



## ORIGINAL ARTICLE

## National estimate and associated factors of myopia among schoolchildren in Ethiopia: A systematic review and meta-analysis

Molla Yigzaw Birhanu<sup>a,\*</sup>, Getamesay Molla Bekele<sup>b</sup>, Cheru Tesema Leshargie<sup>a,c</sup>,  
Selamawit Shita Jemberie<sup>d</sup>

<sup>a</sup> Department of Public Health, College of Health Sciences, Debre Markos university, Debre Markos, Ethiopia

<sup>b</sup> Department of Gynecology and Obstetric, School of Medicine, Debre Markos University, Debre Markos, Ethiopia

<sup>c</sup> Department of Public health, School of Health Sciences, University of Technology Science, Sydney, Australia

<sup>d</sup> Department of Midwifery, College of Health Sciences, Debre Markos University, Debre Markos, Ethiopia

Received 12 August 2022; accepted 25 June 2023

Available online 16 November 2023

### KEYWORDS

Myopia;  
National estimate;  
Associated factors;  
Schoolchildren;  
Ethiopia

### Abstract

**Introduction:** Myopia is a global public health concern that has a significant socioeconomic and psychological impact on schoolchildren. If Myopic patients are not detected early, they are exposed to retinal detachment, cataracts, and glaucoma. There have been previous studies conducted in Ethiopia, but there is significant inconsistency among studies. Hence, the aim of this study was to provide a single figure as well as associated factors for Myopia among Ethiopian schoolchildren.

**Method:** The national and international databases and gray literature were searched for important research articles. This review included school-based cross-sectional studies that were reported in English. The data were extracted using Microsoft word and exported to Stata™ Version 17.0 statistical software for further management and analysis. The presence of heterogeneity was checked using Cochrane Q test via fixed effects model and presented by forest plots with 95% CI. Due to the presence of substantial heterogeneity,  $I^2$  test using random effects model was computed to estimate the effect size. The existing heterogeneity among studies was explained by regional difference. To identify factors associated with myopia, meta regression was computed and significant factors was reported using OR with 95% CI.

**Results:** In this systematic review and meta-analysis, 12 studies with a total of 9688 schoolchildren were included. The national estimate of myopia among schoolchildren in Ethiopia was 6.49% (95%CI: 4.86, 8.12). Having family history of myopia (OR: 9.18, 95%CI: 3.5,24.02) and being female (OR: 0.94, 95% CI: 0.50, 0.98) were the identified factors associated with myopia.

**Conclusion:** Myopia is one of the most prevalent childhood health condition in Ethiopia, which affects about one in every fourteen schoolchildren. Schoolchildren who had family history of myopia and being female were the identified risk factors of myopia among

\* Corresponding author.

E-mail address: [molla\\_yigzaw@dmu.edu.et](mailto:molla_yigzaw@dmu.edu.et) (M.Y. Birhanu).

schoolchildren. Clinical and public engagement activities are needed to address the burden of myopia.

Published by Elsevier España, S.L.U. on behalf of Spanish General Council of Optometry. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## Introduction

Myopia is a visual impairment (refractive error) when accommodation is relaxed, parallel light rays from an object at infinity focus in front of the retina.<sup>1</sup> Because unaided near visual acuity of the myopic eye is usually normal or better than unaided distance acuity (blurry vision), it is commonly referred to as “nearsightedness.”<sup>2</sup> Nearsighted people are more likely to develop vision-threatening conditions such as retinal detachment, choroidal atrophy, cataracts, and glaucoma.<sup>3</sup> Bifocal spectacles, cycloplegic drops, intraocular pressure-lowering drugs, muscarinic receptor antagonists, and contact lenses have all been investigated as interventions to slow the progression of myopia.<sup>4</sup> Almost all patients with myopia can have good vision if they receive early and appropriate correction.<sup>5,6</sup> The burdens of uncorrected myopia are serious especially in students. Poor vision and an inability to read material on the chalkboard due to myopia can greatly affect a student’s participation in education, different activities, and other social participations. Furthermore, when they become adults they are excluded from productive working lives, which results in severe economic and social consequences.<sup>7</sup> Despite the fact that several primary studies have shown that the prevalence of myopia among schoolchildren is high and that prevention and treatment are underutilized, their findings have revealed significant variation in the prevalence of myopia in Ethiopia. The goal of this systematic review and meta-analysis was to determine national estimate and associated factors of myopia among schoolchildren in Ethiopia. The findings of this study, will lay the scientific groundwork for a better understanding of the national estimate of myopia in schoolchildren and which may help to develop effective prevention methods.

Research question: what is the national estimate and associated factors of Myopia among Ethiopian schoolchildren?

Condition: Myopia

Context: Ethiopia

Population: All Ethiopian schoolchildren

## Methods

### Data source and searches

The Preferred Reporting Items for Systematic Review and Meta-Analysis (PRISMA) guideline was used to report this review<sup>8</sup> (Supplementary file1). The five major databases (PubMed/MEDLINE, CINAHL, EMBASE, Google Scholar, and Science Direct) were searched for relevant studies. Moreover, articles were searched from the reference list of eligible studies. The search was carried out independently by three authors (MYB, GMB, and SSJ). Endnote X9 was used to

retrieve and manage studies found through a systematic search. The following terms were used in the above-mentioned databases searches: (((((children[MeSH Terms]) OR (preschool[MeSH Terms])) OR (children[Text Word])) OR (preschool[Text Word]) AND ((y\_1[Filter] AND (fha[Filter]))) AND ((Ethiopia[MeSH Terms]) OR (Ethiopia[Text Word]) AND ((y\_1[Filter] AND (fha[Filter])))) AND (((myopia[Text Word]) OR (refractive error[Text Word])) OR (Myopia[MeSH Terms]) AND ((y\_1[Filter] AND (fha[Filter])))) Filters: Abstract, in the last 1 year.” The searching strategy was lasting from June 28 upto 25 July 2022.

### Eligibility criteria

#### Inclusion criteria

All research articles conducted on myopia among Ethiopian schoolchildren.

#### Study selection procedure

All school based cross-sectional studies were included. All titles/abstracts and full texts found in electronic databases were independently screened by two authors (MYB and GMB). Disagreements were settled through discussion in the presence of third author (CTL) (Fig. 1).

#### Data extraction process and quality assessment

The data were extracted using a Microsoft Excel spreadsheet. To ensure consistency, two authors (MYB and SSJ) extracted data independently using a prepared Microsoft Excel spreadsheet format. During data extraction, the disagreements between data extractor authors were resolved through discussion in the presence of third author (GMB). The data extraction format contains the number of schoolchildren with myopia, the prevalence of myopia in schoolchildren, region, study period, year of publication, sample size, and first author of the primary article to estimate the pooled prevalence of myopia in Ethiopian schoolchildren. As well as, the data extraction checklist contains for factors associated (OR, CI, exposed with disease, nonexposed with nondiseased, exposed with nondiseased, and nonexposed with diseased) with myopia too.

The Newcastle-Ottawa Quality Assessment Scale (NOQAS)<sup>9</sup> was used to evaluate the quality of the included primary studies based on study representativeness, adequate sample size, acceptable non-response rate, use of validated measurement tool, comparability of the study, description of outcome assessment, and use of appropriate statistical tests as parameters. High-quality articles received a global rating of seven out of ten<sup>10,11</sup> (Supplementary file 2).




Fig. 1 PRISMA flow diagram of myopia among Ethiopian schoolchildren.

## Outcome variable

In this systematic review and meta-analysis, national estimate of myopia was the main outcome of interest. The second outcome of this systematic review and meta-analysis was to identify factors associated with myopia among schoolchildren in Ethiopia.

## Heterogeneity, reporting bias, and chance

The presence of heterogeneity among studies was checked using Cochran's-Q test and quantified using I-square statistics. Accordingly, heterogeneity was classified as low, moderate, or high when the values of I-square were 25, 50, and 75%, respectively.<sup>19</sup> Additionally, the dispersion of individual results in the forest plot was also used to evaluate the presence of heterogeneity visually. Egger's linear regression test at a p-value < 0.05 was used to assess the presence of publication bias.<sup>20</sup>

## Data management and analysis

The extracted data were exported to Stata™ Version 17.0 software for further management and analysis. The meta-prop stata command was used to compute the overall pooled estimate. Using a binomial distribution assumption, the standard errors were calculated from the reported estimates and population denominators. The DerSimonian and Laird Method<sup>12</sup> was used to create a random effects meta-analysis model. To identify the possible sources of

heterogeneity, additional statistical analyses such as subgroup analyses, publication bias, and meta-regression were performed. We conducted a subgroup analysis using geographical regions, sample size (above mean Vs. below the mean (807)), and study period (before 2015 Vs. after 2015). Furthermore, sensitivity analysis was performed using a random effects model to assess the impact of a single study on the overall pooled estimate. Finally, the results were presented in tables and forest plots.

## Results

### Search results

A total of 589 studies were found through electronic searching of database such as PubMed, EMBASE, Google Scholar, and Science Direct and others like organizational records and website searches from. About 942 articles were excluded due to duplication, 493 articles were excluded due to different in study context,<sup>13-17</sup> 10 articles were excluded due to different in interest of outcome<sup>18-24</sup> and 5 articles were excluded due to different in study population.<sup>25,26</sup> Finally, 12 cross-sectional studies were identified as having the potential to be included in the current systematic review and meta-analysis.

### Characteristics of included studies

According to Table 1, about 12 studies with a total of 9688 primary and secondary schoolchildren met the inclusion

**Table 1** the characteristics of the studies included in myopia among Ethiopian schoolchildren.

Sn	Authors	Publication year	Region	Study area	Sample size	Prevalence
1	Sara Abebaw <i>et al.</i> <sup>35</sup>	2020	SNNP	Wolkite	1271	7.7
2	Assefa Wolde Yared <i>et al.</i> <sup>30</sup>	2012	Amhara	Gondar	1852	3.5
3	Elias Abera Gebru <i>et al.</i> <sup>33</sup>	2022	SNNP	Hawassa	349	16.05
4	Yemisrach Hailu <i>et al.</i> <sup>37</sup>	2020	Addis Ababa	Addis Ababa	816	2.5
5	Zelalem Addisu Mehari <i>et al.</i> <sup>21</sup>	2013	SNNP	Sodo	4238	6
6	Nebiyat kassa <i>et al.</i> <sup>34</sup>	2014	Addis Ababa	Addis Ababa	1800	2.1
7	Gizachew Tilahun Belete <i>et al.</i> <sup>31</sup>	2017	Amhara	Gondar	498	11.9
8	Jafer Kedir <i>et al.</i> <sup>34</sup>	2014	SNNP	Goro District	570	2.6
9	Tarikayehu Hailemariam <sup>29</sup>	2019	Amhara	Debre Birhan	365	7.8
10	Abel Sinshaw Assem <i>et al.</i> <sup>28</sup>	2021	Amhara	Bahir dara	601	8.49
11	Sintayehu Aweke Sewunet <i>et al.</i> <sup>36</sup>	2014	Amhara	Debre Markos	432	5.47
12	Tibebu Kassa <sup>32</sup>	2003	Amhara	Debark	1134	8.45

criteria. This systematic review and meta-analysis included two regions and one city administration. Among them were Amhara region including six studies<sup>27-32</sup> and South nation and nationality of people ( $n = 4$ ),<sup>21,33-35</sup> and Addis Abeba ( $n = 2$ ).<sup>36,37</sup> The study with the smallest and largest sample sizes, 349 and 4238, was conducted in the SNNPR's Hawassa and Sodo, respectively (Table 1).


**The national estimat of myopia**

In Ethiopia, the national estimat of myopia among schoolchildren was 6.49 percent (95%CI: 4.86, 8.12) (Fig. 2). When

we looked at it by region, Amhara and South nation and nationality of people had largest (7.48 for both of them) and Addis Abeba city administration had smallest burden (2.21% (95%CI:1.65, 2.77)) of myopia (Fig. 3).


**Subgroup meta –analysis**

After determining the presence of heterogeneity between studies, we attempted to address the heterogeneity by conducting subgroup meta-analysis while taking into account region, sample size, and study period. As a result of the subgroup meta-analysis, it was discovered that publication bias



**Fig. 2** The overall pooled prevalence of myopia among schoolchildren in Ethiopia.

## The regional pooled prevalence of Myopia in Ethiopia



**Fig. 3** The regional pooled prevalence of myopia in Ethiopia.

study period, and sample size were all identified as sources of heterogeneity with the presence of strong evidence supporting the existence of heterogeneity that means region (Amhara:  $I^2=92.2$  with p-value 0.001, SNNP:  $I^2=94.7$  with p-value 0.001, and Addis Ababa:  $I^2 = 0.0$  with p-value 0.53) (Fig. 4), studies having sample size  $> 807$ :  $I^2 = 96.00$  with p-value 0.0001 and those studies having sample size  $< 807$ :  $I^2= 93.4$  with p-value 0.0001 (Fig. 5), study period(after 2015:  $I^2= 93.4$  with p-value 0.0001 and before 2015:  $I^2= 93.4$  with p-value 0.0001) (Fig. 6).

### Meta-regression

The publication year and sample size were used as covariates in random-effects meta-regression. The analysis revealed that sample size ( $p = 0.72$ ) and publication year ( $p = 0.99$ ) had no effect on heterogeneity (heterogeneity was not explained by sample size and publication year) (Table 2).

### Publication bias (bias detection)


A funnel plot was used to determine the presence or absence of publication bias, and the scatter plot points were

asymmetrical (attributed to publication bias), indicating that the small-study had effects on the heterogeneity of myopia estimate among Ethiopian schoolchildren (Fig. 7). The Egger linear regression test was used to objectively check the presence or absence of publication bias. As a result, there was statistically significant publication bias ( $P = 0.016$ ) (Table 3).

### Factors associated with myopia

Factors associated with myopia in schoolchildren were determined based on sex (including seven studies),<sup>27-29,31-33,37</sup> family history of myopia (including four studies),<sup>28,29,31,33</sup> residence(including eight studies),<sup>21,28-31,34,36,37</sup> family educational status (including four studies)[29–31, 34], and use of computer regularly (three studies).<sup>28,35,37</sup> Finally, being female and having family history of myopia were the factors which are significantly associated with myopia.

Being rural residence was 1.09 (95% CI: 0.56, 2.13), had no regular computer exposure had a 0.69 (95% CI: 0.45, 1.06), and having educated family had a 1.89 (95% CI: 0.59, 6.06) times larger chance of myopia than its corresponding counterpart. However, these differences were not statistically significant.



**Fig. 4** Subgroup analysis using sample size, 807 as cut off point.

The odds of myopia among female schoolchildren was 0.94 (95% CI: 0.50, 0.980) times lower than male schoolchildren (Fig. 6). Schoolchildren who had family history of myopia was 6.48 (95% CI: 3.03, 13.82) times greater than students who had no family history of myopia (Fig. 8).

## Discussions

This systematic review and meta-analysis incorporated 12 papers with 9688 research subjects. These studies were conducted between 2003 and 2021. The subjects were selected using either a simple random or systematic random sampling approach in each study. The sample size for included studies were ranged from 349 to 4238 individuals. The participated children were ranged from 7 to 15 years old with the median age of  $10 \pm 2$  interquartile range. The spherical equivalent was considered in measuring myopia including cycloplegia and it was measured from  $-1.5$  diopters to  $-6$  diopters.

The overall national estimate of myopia among schoolchildren in Ethiopia was 6.49% (95%CI: 4.85, 8.12). This study's findings are consistent with the study's findings that the overall estimated prevalence in East Africa was 4.9% in 2010 and 8.4% in 2020.<sup>38</sup>

It is supported by a study carried out in India (7.5%).<sup>39</sup> The finding of this systematic review and meta-analysis is higher than the finding of the systematic review and meta-analysis conducted in Africa (4.7%) among schoolchildren.<sup>40</sup> This could be due to the fact that the parents in this study had less educational exposure than the previous one. Educated families are more concerned than their counterparts about their children's early screening as well as diagnosis of myopia for the early initiation of the treatment. This could explain the higher prevalence of myopia among schoolchildren in our study.

The prevalence found in this study was lower than the study finding conducted in Nigeria where it was found a 33.9%<sup>41</sup> and Chidambaram, Tamil Nadu India 12. This could be because of the study participants in this study included both primary and secondary schoolchildren, whereas the previous one included only primary schoolchildren. Myopia is more common in primary schoolchildren than among secondary schoolchildren.<sup>42</sup> In our study, there was decrease of the infected schoolchildren in the presence of larger denominator which can decrease the burden of the myopia than their counterparts. This could be the possible reason for the higher prevalence of myopia in the previous study than the present one.

The odds of experiencing myopia among schoolchildren who had family history of myopia were 6.48 (95% CI: 3.03,




Fig. 5 Subgroup meta-analysis using study period after and before 2015.





Fig. 6 s: Funnel plot to check publication bias.

13.82) times greater than their counterparts. This is consistent with the study conducted in Cardiff University.<sup>43</sup> This could mean that children born from myopic families are more likely to develop myopia themselves.<sup>44</sup> This could be one of the reasons why schoolchildren with a family history of myopia have a higher risk of developing myopia than their peers.

**Limitation of the study**

This study’s national estimate of myopia may not exactly represent the burden of Ethiopian schoolchildren because it was conducted using primary studies only from Amhara, South nation nationalities and people, and Addis Ababa. In addition to the foregoing, this study only included articles published in English, which may affect the prevalence of myopia and taken as a limitation of this study.


Logrr	Coefficient	Std. err.	t	P>  t	[95% conf. interval]
Sample size	< 0.0001	0 0.0002	0.37	0.72	−0.001, 0 0.001
Publication year	< −0.0001	0.005	−0.02	0.99	−0.12, 0.120
Constant	3.03	109.47	0.03	0.98	−24.13, 25.68



**Fig. 7** Association between myopia and family history of myopia in Ethiopia.

**Table 3** The table to check the publication bias objectively.

Std_Eff	Coefficient	Std. err.	T	P>  t	[95% conf. interval]
Slope	1.26	1.24	1.01	0.34	-1.51, 4.02
Bias	6.01	2.07	2.90	0.016	1.39, 10.62



**Fig. 8** Association between myopia and family history of myopia in Ethiopia.

## Conclusion

One of the most prevalent childhood health condition in Ethiopia is myopia, which affects about one in every fourteen schoolchildren. Having family history of myopia was the identified risk factors of myopia. Clinical and public engagement activities are needed to address the burden of myopia. Public engagement activities like providing health education for parents and students to have early screening and diagnosis of myopia whether they present with myopia or not and clinical engagement activities like early diagnosis and treatment of myopia need to be given great emphasis.

## Recommendation

The schoolchildren who were from parents having myopia should check their visual acuity regularly. Parents and schools should inform and remind the students to check their visual acuity regularly and take the correction if they develop the disease already. The health institutions should provide health education regarding myopia for students, parents, and schools about the prevention and control measures of myopia.

## Contributors

Conceptualization: MYB, SSJ  
 Data curation: MYB, GMB, SSJ.  
 Formal analysis: MYB, CTL, SSJ.  
 Investigation: MYB, CTL, SSJ.  
 Methodology: MYB, GMB, CTL, SSJ.  
 Project administration: MYB, CTL, SSJ.  
 Resources: MYB, CTL, SSJ.  
 Software: MYB, CTL, GMB, SSJ.  
 Supervision: MYB, SSJ.  
 Validation: MYB, CTL, SSJ, GMB.  
 Visualization: MYB, CTL, SSJ  
 Writing – original draft: MYB, GMB, CTL, SSJ.  
 Writing – review & editing: MYB, GMB, CTL, SSJ.

## Ethics consideration

N/A.

## Funding

N/A.

## Conflicts of interest

The authors declared that they didn't have any form of competing interest.

## Supplementary materials

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.optom.2023.100480](https://doi.org/10.1016/j.optom.2023.100480).

## References

- Goss DA, Grosvenor TP, Keller JT, Marsh-Tootle W, Norton TT, Zadnik K. Optometric clinical practice guideline care of the patient with myopia. *Am Optometr Assoc.* 1997;243:63141–67881.
- Wensor M, McCarty CA, Taylor HR. Prevalence and risk factors of myopia in Victoria, Australia. *Arch Ophthalmol.* 1999;117(5):658–663.
- Jin J-X, Hua W-J, Jiang X, et al. Effect of outdoor activity on myopia onset and progression in school-aged children in north-east China: the Sujiatun Eye Care Study. *BMC Ophthalmol.* 2015;15(1):1–11.
- Daoud YJ, Hutchinson A, Wallace DK, Song J, Kim T. Refractive surgery in children: treatment options, outcomes, and contraindications. *Am. J. Ophthalmol.* 2009;147(4):573–582. e572.
- Wu LJ, You QS, Duan JL, et al. Prevalence and associated factors of myopia in high-school students in Beijing. *PLoS One.* 2015;10(3): e0120764.
- Pan CW, Ramamurthy D, Saw SM. Worldwide prevalence and risk factors for myopia. *Ophthalmic Physiol Opt.* 2012;32(1):3–16.
- Congdon N, Wang Y, Song Y, et al. Visual disability, visual function, and myopia among rural Chinese secondary school children: the Xichang Pediatric Refractive Error Study (X-PRES)—report 1. *Invest. Ophthalmol. Vis. Sci.* 2008;49(7):2888–2894.
- Rethlefsen ML, Kirtley S, Waffenschmidt S, et al. PRISMA-S: an extension to the PRISMA statement for reporting literature searches in systematic reviews. *Syst Rev.* 2021;10(1):1–19.
- Fazel S, Khosla V, Doll H, Geddes J. The prevalence of mental disorders among the homeless in western countries: systematic review and meta-regression analysis. *PLoS Med.* 2008;5(12): e225.
- Islam MM, Iqbal U, Walther B, et al. Benzodiazepine use and risk of dementia in the elderly population: a systematic review and meta-analysis. *Neuroepidemiology.* 2016;47(3–4):181–191.
- Leshargie CT, Alebel A, Kibret GD, et al. The impact of peer pressure on cigarette smoking among high school and university students in Ethiopia: a systemic review and meta-analysis. *PLoS One.* 2019;14(10): e0222572.
- DerSimonian R, Laird N. Meta-analysis in clinical trials. *Control Clin Trials.* 1986;7(3):177–188.
- Atowa UC, Wajuihian SO, Munsamy AJ. Prevalence and risk factors for myopia among school children in Aba, Nigeria. *Afr Vis Eye Health.* 2017;76(1):1–5.
- Atowa UC, Wajuihian SO, Munsamy AJ. Associations between near work, outdoor activity, parental myopia and myopia among school children in Aba, Nigeria. *Int J Ophthalmol.* 2020;13(2):309.
- Uhuama O, Atowa U. Distribution of refractive errors among school children in Abia state of Nigeria. *J Nigerian Optomet Assoc.* 2004:11.
- Ovenseri-Ogbomo G, Omuemu V. Prevalence of refractive error among school children in the Cape Coast Municipality, Ghana. *Clin Optom.* 2010;2:59–66.
- Ovenseri-Ogbomo G, Assien R. Refractive error in school children in Agona Swedru, Ghana. *Afr Vis Eye Health.* 2010;69(2):86–92.
- Ajaiyeoba AI, Isawumi MA, Adeoye AO, Oluleye TS. Pattern of eye diseases and visual impairment among students in south-western Nigeria. *Int Ophthalmol.* 2007;27(5):287–292.

19. Megbelayin O, Asana E. Visual impairment among schoolchildren calabar vision screening survey in secondary schools (CVS4 study). *Internet J Ophthalmol Visual Sci.* 2013;10(1):1–7.
20. Alabi AS, Aribaba OT, Alabi AO, Ilo O, Onakoya AO, Akinsola FB. Visual impairment and ocular morbidities among schoolchildren in Southwest, Nigeria. *Nigerian Postgraduate Med J.* 2018;25(3):166.
21. Mehari ZA, Yimer AW. Prevalence of refractive errors among schoolchildren in rural central Ethiopia. *Clin Exp Optometry.* 2013;96(1):65–69.
22. Nebiyat K, Alemayehu W, Tigist S. Refractive errors among school children in Addis Ababa, Ethiopia. *J Ophthalmol Eastern, Cent Southern Afr.* 2015;19(2).
23. Zelalem M, Abebe Y, Adamu Y, Getinet T. Prevalence of visual impairment among school children in three primary schools of Sekela Woreda, Amhara regional state, north-west Ethiopia. *SAGE Open Med.* 2019;7: 2050312119849769.
24. Dhanesha U, Polack S, Bastawrous A, Banks LM. Prevalence and causes of visual impairment among schoolchildren in Mekelle, Ethiopia. *Cogent Med.* 2018;5(1): 1554832.
25. Assefa NL, Admas AW, Adimasu NF. Prevalence and associated factors of visual impairment among adults at Debre Berhan town, North Shewa, Ethiopia. *BMC Ophthalmol.* 2020;20(1):1–8.
26. Alemu DS, Gudeta AD, Ferede AT, Alemu HW. Prevalence and degrees of myopia and hyperopia at Gondar university hospital tertiary eye care and training center, Northwest Ethiopia. *Clin Optometry.* 2016;8:85.
27. Sewunet SA, Aredo KK, Gedefew M. Uncorrected refractive error and associated factors among primary school children in Debre Markos District, Northwest Ethiopia. *BMC Ophthalmol.* 2014;14(1):1–6.
28. Assem AS, Tegegne MM, Fekadu SA. Prevalence and associated factors of myopia among school children in Bahir Dar city, Northwest Ethiopia, 2019. *PLoS One.* 2021;16(3): e0248936.
29. Tarikayehu H.: Prevalence of myopia and associated factor among elementary school children in Debre Berhan Town Amhara Regional State, North Shoa Ethiopia. 2019.
30. Yared AW, Belaynew WT, Destaye S, Ayanaw T, Zelalem E. Prevalence of refractive errors among school children in Gondar town, northwest Ethiopia. *Middle East Afr J Ophthalmol.* 2012;19(4):372.
31. Belete GT, Anbesse DH, Tsegaye AT, Hussen MS. Prevalence and associated factors of myopia among high school students in Gondar town, northwest Ethiopia. *Clin Optometry.* 2016;9:11.. 2017.
32. Kassa T, Alene GD. Prevalence of refractive errors in pre-school and school children of Debarke and Kola Diba towns, North-western Ethiopia. *Ethiopian J Health Dev.* 2003;17(2):117–124.
33. Gebru EA, Mekonnen KA. Prevalence and factors associated with myopia among high school students in Hawassa City, South Ethiopia, 2019. *Clin Optometry.* 2022;14:35.
34. Kedir J, Girma A. Prevalence of refractive error and visual impairment among rural school-age children of Goro District, Gurage Zone, Ethiopia. *Ethiop J Health Sci.* 2014;24(4): 353–358.
35. Gessesse SA, Teshome AW. Prevalence of myopia among secondary school students in Welkite town: South-Western Ethiopia. *BMC Ophthalmol.* 2020;20(1):1–6.
36. Woldeyes A, Misganaw C. prevalence and factors associated with refractive error among primary school children in Addis Ababa, Ethiopia. *Int J Med Health Sci Res.* 2014;1 (9):92–104.
37. Hailu Y, Hiko D, Shaweno T. Prevalence of visual impairment and associated factors among primary schoolchildren in Addis Ababa, central Ethiopia. *Clin Ophthalmol.* 2020;14:767.
38. Holden BA, Fricke TR, Wilson DA, et al. Global prevalence of myopia and high myopia and temporal trends from 2000 through 2050. *Ophthalmology.* 2016;123(5):1036–1042.
39. Agarwal D, Saxena R, Gupta V, et al. Prevalence of myopia in Indian school children: meta-analysis of last four decades. *PLoS One.* 2020;15(10): e0240750.
40. Oveneri-Ogbomo G, Osuagwu UL, Ekpenyong BN, et al. Systematic review and meta-analysis of myopia prevalence in African school children. *PLoS One.* 2022;17(2): e0263335.
41. Xie Z, Long Y, Wang J, Li Q, Zhang Q. Prevalence of myopia and associated risk factors among primary students in Chongqing: multilevel modeling. *BMC Ophthalmol.* 2020;20(1):1–8.
42. Guo L, Yang J, Mai J, Du X, Guo Y, Li P. *Prevalence and Associated Factors of Myopia Among Primary and Middle School-Aged Students: A School-Based Study in Guangzhou.* 2016.
43. Farbrother JE, GK, Owen MJ, Guggenheim JA. *Family Aggregation of High Myopia: Estimation of the Sibling Recurrence Risk Ratio.* 2004.
44. Mu MHe MM. *The Effect of Having Myopic Parents: An Analysis of Myopia in Three Generations.* 1999.