



LETTER TO THE EDITOR

On the urgency of air pollution control to manage chronic eye rubbing and probable risk of keratoconus



Dear editor in chief,

This letter attempts to address the linkage between air pollution and ocular allergies leading to other complications such as keratoconus (KCN) disease. The rapid growth of urbanization and continued industry development have led to the rising emission of many different pollutants into ambient air. Also, the heating and cooling of modern houses are gaining importance to reduce costs; accordingly, recent buildings are facing poor air quality due to air-tight and poor ventilation conditions. Current literature shows that exposure to air pollution has made the greatest hazard to major body systems such as the respiratory and cardiovascular systems. Also, healthcare costs and hospital admissions are enhanced as this phenomenon escalates. Overall, it is confirmed that the health concerns associated with air pollution tend to worsen with time around the world.^{1,2}

In recent decades, the prevalence of allergic reactions has constantly grown worldwide and air pollution is considered a major contributing factor. Allergies may trigger various symptoms such as conjunctivitis, dry eyes, runny nose, sneezing, coughing, itchy throat, eczema, and certain inflammatory skin disease. It is worth noting that air pollutants such as dust, NO₂, O₃, PM_{2.5}, PM₁₀, SO₂, CO, pollen, dirt, and lint either indoors or outdoors play the greatest role in the incidence of allergy symptoms. These airborne pollutants are known as important sources of air pollutants that originate from plenty of natural and anthropogenic sources, such as vehicle exhaust, power plants, fossil fuels, volcanoes, forest fires, waste deposits, agriculture, atmospheric photochemical reactions, cooking stoves, and heating systems.³

Primary care providers often overlook ophthalmological complaints arising from allergic responses. These ocular complaints mostly include dryness, irritation, redness, foreign body sensation, tears, and blurring of vision. The starting point for treatment of these symptoms includes the use of tear substitutes, which flush allergens from the ocular surface, as well as topical antihistamines and mast cell stabilizers. Corticosteroids remain among the most potent pharmacologic agents used in the more severe variants of

ocular allergy and are effective in the treatment of acute and chronic forms of ocular allergic conjunctivitis. While physicians who currently treat ocular allergy have a greater familiarity with the signs and symptoms of allergic disease, misuse of topical and systemic corticosteroids by patients may even cause severe complications of acquired cataracts and secondary glaucoma.⁴ It is reported that air pollution exposure exacerbates allergic symptoms in the eye through oxidative stress, toxicity, and inflammation.² Consequently, unaddressed allergic eye symptoms trigger the habit of chronic eye rubbing; Rubbing can worsen tissue damage and increase proteases like matrix metalloproteinase-9. Elevated serum IgE indicates an allergy, triggering histamine and IL-13 release, which activates itch receptors.⁵ Unfortunately, chronic eye rubbing may lead to the development and progression of keratoconus or KCN disease. 48% of patients with keratoconus have eye-rubbing behavior ($P = 0.018$).⁶ Through an indirect association, such complications are likely to develop during vigorous and prolonged rubbing habits.⁷

The KCN disease is a non-inflammatory disease of the cornea, where the cornea thins and changes its shape. Originally, the cornea is round or spherical, but during keratoconus, the cornea becomes convex and conical. Thinning and changing the shape of the cornea from spherical to conical form leads to irregular myopia-astigmatism and thus reduces vision. Although the etiology of KCN is complex and involves a multifactorial origin such as demographic and environmental factors, some studies confirm the role of eye rubbing as an important risk factor in developing keratoconus.⁸

Therefore, air pollutants both directly and indirectly constitute an imminent risk for the occurrence of keratoconus. In direct pathways, they are thought to exacerbate known risk factors such as eye rubbing and atopy. In an indirect pathway, air pollutants like PM could affect the cornea by inducing the apoptosis of epithelial cell types and interacting directly with stromal collagen.⁹ studies shows a significant positive correlation between prevalence of keratoconus with particles rate of PM_{2.5} ($R = 0.58$; $P < 0.001$), PM₁₀ ($R = 0.67$; $P < 0.001$) and NO₂ ($R = 0.64$; $P = 0.00016$).¹⁰

To conclude, we would like to close this letter by warning that air pollution through allergic development and subsequently atopy and eye rubbing should be introduced as an important risk factor in developing keratoconus in the

<https://doi.org/10.1016/j.optom.2025.100554>

1888-4296/© 2025 The Authors. Published by Elsevier España, S.L.U. on behalf of Spanish General Council of Optometry. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

population exposed. The increased incidence of keratoconus further underscores the importance of air pollution control and prevention of individual exposure to preventable risks through effective policies and strategies as part of the responsibility of the governments. This letter also highlights the need to enhance public awareness about the effect of eye rubbing on keratoconus and the necessity of treatment and prevention of itching and dryness as the main causes of eye rubbing.

CRedit authorship contribution statement

Ebrahim Shirzadeh: Conceptualization. **Nematullah Shomoossi:** Writing – original draft. **Hadi Hasani:** Writing – review & editing, Project administration, Supervision.

References

1. Mandell JT, Idarraga M, Kumar N, Galor A. Impact of air pollution and weather on dry eye. *J Clin Med*. 2020;9(11):3740. <https://doi.org/10.1016/j.jtos.2018.03.001>.
2. Jung SJ, Mehta JS, Tong L. Effects of environment pollution on the ocular surface. *Ocul Surf*. 2018;16(2):198–205. <https://doi.org/10.1016/j.jtos.2018.03.001>.
3. Kim K-H, Jahan SA, Kabir E. A review of human health perspective of air pollution with respect to allergies and asthma. *Environ Int*. 2013;59:41–52. <https://doi.org/10.1016/j.envint.2013.05.007>.
4. Shirzadeh E, Shomoossi N, Abdolazadeh H. Topical anesthetic misuse in patients admitted to Sabzevar Eye Clinic in Iran. *Acta facultatis medicae Naissensis*. 2016;33(4):287–294.
5. Shetty R, Sureka S, Kusumgar P, Sethu S, Sainani K. Allergen-specific exposure associated with high immunoglobulin E and eye rubbing predisposes to progression of keratoconus. *Ind J Ophthalmol*. 2017;65(5):399–402. https://doi.org/10.4103/ijo.IJO_217_17.5.
6. Weed KH, MacEwen CJ, Giles T, et al. The Dundee University Scottish Keratoconus study: demographics, corneal signs, associated diseases, and eye rubbing. *Eye (Lond)*. 2008;22:534–541.
7. Ben-Eli H, Erdinest N, Solomon A. Pathogenesis and complications of chronic eye rubbing in ocular allergy. *Curr Opin Allergy Clin Immunol*. 2019;19(5):526–534. <https://doi.org/10.1097/aci.0000000000000571>.
8. Najmi H, Mobarki Y, Mania K, et al. The correlation between keratoconus and eye rubbing: a review. *Int J Ophthalmol*. 2019;12(11):1775–1781. <https://doi.org/10.18240/ijo.2019.11.17>.
9. Jurkiewicz T, Marty A-S. Correlation between keratoconus and pollution. *Ophthalmic Epidemiol*. 2021;28(6):495–501.
10. Jurkiewicz T, Marty AS. Air pollution and the prevalence of keratoconus: is there a connection? *Ophthalmic Epidemiol*. 2024;1–9. <https://doi.org/10.1080/09286586.2024.2399765>.

Ebrahim Shirzadeh^a, Nematullah Shomoossi^b, Hadi Hasani^{c,*}

^a Department of Ophthalmology, School of Medicine, Sabzevar University of Medical Sciences, Sabzevar, Iran

^b School of Medicine, Sabzevar University of Medical Sciences, Sabzevar, Iran

^c Department of Medical Surgical Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran

* Corresponding author at: Department of Medical Surgical Nursing, School of Nursing and Midwifery, Tehran University of Medical Sciences, Tehran, Iran.

E-mail addresses: hasani1376@gmail.com, h-hasani@razi.tums.ac.ir (H. Hasani).