



REVIEW

Evidence on the parameters of oculomotor skills and normative values: A systematic review



Mario Cantó-Cerdán^{a,b,*}, Carlos Javier Hernández-Rodríguez^{c,d}, Antonio Martínez-Abad^{a,b}

^a Department of Optometry, Vissum (Miranza Group), Alicante, Spain

^b Grupo de Investigación en Optometría (GIOptom), University of Alicante, Alicante, Spain

^c Department of Optometry, OKYO Ophthalmics, Alicante, Spain

^d Department of Ophthalmology, Quirónsalud Medical Center, Elche, Spain

Received 24 March 2025; accepted 14 June 2025

Available online 10 July 2025

KEY WORDS

Oculomotor skills;
Oculomotor function;
Systematic review;
Gaze processing;
Saccadic

Abstract

Purpose: To evaluate the current evidence on oculomotor measurement parameters and their normative values through a systematic review.

Methods: A search of primary studies was conducted using a search equation with free language. Original articles analyzing normal oculomotor function parameters in healthy populations of any age, studies that included a clearly differentiated healthy control group, and articles using any oculomotor measurement test were included. The QUADAS-2 tool was used to assess the risk of bias, applicability, and quality of the studies. The review was conducted independently by the authors and then pooled to determine the final inclusion.

Results: A total of 915 articles were identified, of which 750 were excluded after the first review of the title and abstract. In the second step, 133 out of 165 investigations were discarded. Ultimately, 32 articles from the initial search were included, along with 10 additional articles identified through a manual search. The findings revealed variations in how oculomotor skills are measured, including differences in stimuli, measurement distances, and parameters assessed. A high risk of bias was observed ($\geq 50\%$ in the areas of “flow and timing”, “reference standard” and “patient selection”) along with poor applicability ($\geq 50\%$ in all aspects).

Conclusions: There is no clear evidence on normative values for oculomotor skills, nor is there a consensus on the measurement methods, stimulus used, or working distance. Furthermore, there is no agreement on which aspects of oculomotor skills should be assessed. To enhance reliability and applicability, measurement criteria should be standardized, and normative values should be established.

© 2025 The Authors. Published by Elsevier España, S.L.U. on behalf of Spanish General Council of Optometry. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

* Corresponding author: Mario Cantó-Cerdán, University of Alicante, Avda de Denia s/n, Vissum Alicante 03016 Alicante, Spain.
E-mail address: mario.canto@vissum.com (M. Cantó-Cerdán).

Introduction

Oculomotor function refers to an individual's innate ability to execute eye movements in a seamless, coordinated, and fluid manner, ensuring the maintenance of a clear, fused, and stable image on the central region of the retina. When an object is in motion, it is crucial to sustain this visual stability as the object moves. To achieve accurate oculomotor performance, three fundamental skills must be assessed: tracking movements, saccadic movements, and fixation movements. These skills should be well developed monocularly and must also be coordinated to ensure efficient oculomotor function.¹

Oculomotor skills are currently being studied across various disciplines within the health sciences.² Most research is focused on the neurological field, including schizophrenia,³ cognitive impairment,⁴ depression,⁵ biomarkers of neurodegenerative diseases,⁶ neurodevelopmental disorders,⁷ and even post-COVID conditions.⁸ Additionally, oculomotor skills play a crucial role in non-health-related fields, such as marketing⁹ or enabling individuals with amyotrophic lateral sclerosis to communicate using eye-tracking technology.¹⁰

In the field of optometry, oculomotor studies have grown over the past decade, largely due to advancements in eye-tracking technologies.¹¹ Research has focused on various areas, including visual dysfunctions, strabismus, amblyopia, nystagmus, and visual impairment. Additionally, oculomotor studies have extended to vision-related fields such as refractive surgery, sports vision or driving vision.^{2,11,12}

However, despite this exponential growth, concerns remain regarding the diagnostic methodology for oculomotor abnormalities. No clear range of normal values has been established for the parameters used to assess ocular motility, and no gold standard or reference test has been defined.¹³ Classical tests, such as the Developmental Eye Movement Test (DEM),¹⁴ the Northeastern State University College of Optometry (NSUCO) Oculomotor Test,¹⁵ and the King-Devick Test (K-D)¹⁶ have been used for many years. However, these tests have the disadvantage of being subjective for both the examiner and the patient. With the advent of new eye-tracking technologies, equipment has been developed to measure oculomotor parameters objectively. However, these systems vary in the parameters they assess, the psychophysics of measurement, the stimulus used, and the measurement distance.^{11–13}

Research has been conducted to establish normative ranges for different subjective tests,^{14–16} as well as for some of the new technologies.^{17–20} However, there is still no scientific consensus regarding the psychophysics of measurement, the range of values, or the parameters to be assessed. This lack of standardization highlights the need for a unified approach to these aspects.

Therefore, the aim of this study is to evaluate the current evidence on oculomotor measurement parameters and their normative values through a systematic review.

Methods

The study has been registered in the PROSPERO database. A comprehensive search was conducted using a search strategy ([Supplementary Material](#)) in three databases: PubMed,

Web of Science, and Scopus. The search included all age groups and imposed no time restrictions to ensure that no relevant articles were overlooked. After the initial search, a refined selection of articles was made based on the following criteria:

Inclusion criteria

- Original descriptive articles analyzing normal oculomotor parameters (pursuit, saccadic movements, fixation, saccadic velocity, and reading saccades) in a healthy population of any age.
- Originals comparative articles analyzing oculomotor parameters, whose design includes a well-defined control group not accomplishing inclusion criteria.
- Articles utilizing a subjective test currently used in the clinical practice such as NSUCO, DEM, ADEM, or K-D.
- Articles utilizing an objective test currently used in the clinical practice such as eye-tracker, or video-oculography.

Exclusion criteria

- Case series, clinical case reports, and animal studies.
- Studies involving populations with neurological, developmental, or ocular pathologies that lack a well-defined control group or include fewer than 20 subjects.
- Methods employing stimuli different to standard geometrical shapes (dot, cross, square or similar) or related to language (numbers, letters).
- Methods providing non quantitative results such as graphs or interpretations.
- Samples where visual evaluations were performed diagnosing binocular anomalies, such as amblyopia and strabismus in its subjects.

The article selection process followed a sequential approach. First, titles and abstracts were reviewed to exclude irrelevant studies, and duplicates were removed. Second, full-text articles were examined, and only those that met the predefined inclusion criteria and addressed the research question were selected. Third, a manual search considering studies known by articles and reviewing references of included articles was conducted to identify additional references that might not have appeared in the initial search. Articles with control groups that did not clearly report results were excluded.

Finally, to assess the risk of bias, applicability, and quality of the studies, the QUADAS-2 tool was used. This tool is divided into domains that evaluate key aspects of each article, such as participant selection, index test, reference test, and the flow and timing of the study. Following the guidelines of the evaluation tool, the risk of bias was classified into three evidence level groups: studies with a low risk of bias, studies with an unclear risk of bias, and studies with a high risk of bias.

The article selection and the quality analysis using the QUADAS-2 procedure were performed blindly and independently by three investigators to ensure accurate classification of the included studies. In cases of discrepancies, a consensus was reached between the authors, with all three authors discussing and agreeing. (REF QUADAS-2 2011 Withing)




Fig. 1 Flow chart of eligible papers used in the meta-analysis (PRISMA statement). PRISMA, preferred reporting items for systematic reviews.

The initial literature search was conducted in February 2024, and the databases were reviewed again in September 2024 using the same methodology.

Results

The study selection process for this systematic review is illustrated in a flowchart in Fig. 1. Initially, 915 documents were retrieved during the search. After reviewing titles and abstracts and removing duplicates, 165 articles were selected for full-text evaluation. Of these, 133 were excluded for not meeting the inclusion criteria. Ultimately, 32 articles were included. Additionally, a manual search identified 6 more articles, bringing the total to 38 articles that were analyzed in this review.

Study characteristics

Table 1 presents the characteristics of the 38 included studies, which span from 1980 to 2023. The studies included sample sizes ranging from 20 to 2075 subjects. Of these, 22 studies (57.9 %) were descriptive cross-sectional, 15 studies

(39.5 %) were comparative cross-sectional, and 1 study (2.6 %) was pseudo-experimental.

Regarding oculomotor skills, 34 out of the 38 studies (89.5 %) addressed saccadic movements, 16 studies (42.1 %) focused on tracking movements, and 11 studies (28.9 %) investigated fixation movements. In terms of technology and measurement systems, a significant amount of variability was found, with classical tests such as the DEM test, NSUCO test, and electro-oculography in studies performed >30 years ago, alongside newer technologies, like eye trackers, video-oculography, and video-nystagmography in recent evidence. The stimuli used also varied widely, including letters, texts, reading tests, numbers, and monochromatic, colored, or illuminated Figs. It diffculted the stratified analysis by type test. Similar limitation was found for age because 18 of 38 studies (47.4 %) was performed exclusively in children but ones in babies and others in adolescents, 15 of 38 (39.5 %) were performed exclusively in adults, and 5 of 38 (13.2 %) was performed combining both population. Notably, only 12 of the 38 studies (31.6 %) conducted an optometric examination prior to the oculomotor assessment. Finally, regarding the working distance for measuring oculomotor values, variability was observed, ranging from 25 cm to 200 cm. It is important to note that 9 studies (23.7 %) did not specify the working distance, and 2

Table 1 Characteristics of the 38 studies included in the review.

Year	Author	Characteristics of the control group (n, ranged age)	Study design	Oculomotor skills	Measurments system	Stimulus	Optometric examination	Work Distance (cm)
1980	Schalen ²¹	20, age 22–70	Cross-sectional, descriptive	SAC, SP	Electro-oculography	Light Spot in computer	No	160 cm
1984	Bergenius ²²	60, age 11–70	Cross-sectional, descriptive	SAC, SP	Electro-oculography	Light Spot in computer	No	70 cm
1990	Garzia ¹⁴	534 normal subjects aged 6–13	Cross-sectional, descriptive	SAC	DEM	Numbers on test card	No	33 cm
1992	Maples ¹⁵	1714 normal subjects aged 5–14	Cross-sectional, descriptive	SAC, SP	NSUCO	Sphere	No	Not described
1992	Versino ²³	76 normal subjects aged 12–77	Cross-sectional, descriptive	SAC	Electro-oculography	Light Spot	No	Not described
1993	Ross ²⁴	53 normal subjects aged 7–15	Cross-sectional, descriptive	SAC, SP	Eyetracker	Small target in computer	No	43 cm
1997	Litman ²⁵	24 normal subjects	Cross-sectional, comparative	SAC, SP	Eyetracker	Bright Square	No	43 cm
1997	Ross ²⁶	25 normal subjects	Cross-sectional, comparative	SAC, SP	Eyetracker	White disc in screen	No	120 cm
1999	Campana ²⁷	65 normal subjects	Cross-sectional, comparative	SAC	Eyetracker	Light Spot	No	90 cm
2003	Jimenez ²⁸	1056 normal subjects aged 6–12	Cross-sectional, descriptive	SAC	DEM and Schei-man & Wick procedure	Numbers on test card and fixing point	Yes	40 cm // 33 cm DEM
2006	Rutsche ¹⁸	358 normal subjects aged 0–6	Cross-sectional, descriptive	SAC, SP	Pupil reflex	Color lights	Yes	40 cm
2007	Aring ²⁹	135 normal subjects aged 4–15	Cross-sectional, descriptive	FIX	Infrared light	Dot	Yes	53 cm
2010	van Tritch ³⁰	28 normal subjects aged 15–35	Cross-sectional, comparative	SAC, SP	Double magnetic induction method	Red light laser spot	No	Not described
2011	Goepel ³¹	31 normal subjects age 7–12	Cross-sectional, descriptive	SAC	Eyetracker	Cartoons in computer	No	51,8 cm
2011	Kattoulas ³²	2075 normal subjects aged 18–25	Cross-sectional, descriptive	SAC, SP, FIX	Eyetracker	Cross	No	Not described
2011	Webber ³³	59 normal subjects aged 9–10	Cross-sectional, descriptive	SAC	Eyetracker and DEM	Numbers on test card	Yes	Not described
2012	Shi ³⁴	20 normal subjects aged 6–12	Cross-sectional, comparative	FIX	Eyetracker	Light spot	Yes	200 cm
2012	Boot ³⁵	213 normal subjects aged 0–12	Cross-sectional, descriptive	FIX	Eyetracker	White Dots	No	60 cm

Table 1 (Continued)

Year	Author	Characteristics of the control group (n, ranged age)	Study design	Oculomotor skills	Measurments system	Stimulus	Optometric examination	Work Distance (cm)
2015	Seferlis ³⁶	250 normal subjects aged 18–70	Cross-sectional, descriptive	SAC, SP, FIX	Video-oculography	Light spot	Yes	170 cm
2015	Doettl ³⁷	62 normal subjects: 4–44	Cross-sectional, comparative	SAC, SP	Videonystagmography	Snellen chart	Yes	127 cm
2016	Zalla ³⁸	20 normal subjects	Cross-sectional, comparative	SAC	Eyetracker	Grid of 13 points	No	60 cm
2016	Yang ³⁹	50 normal subjects aged 20–69	Cross-sectional, descriptive	SAC, SP	Video-oculography	Point	No	100 cm
2017	Choi ⁴⁰	63 normal subjects: aged 19–80	Cross-sectional, comparative	SAC, FIX	Eyetracker	Text	No	85 cm
2017	Ferreira ⁴¹	31 normal subjects aged 18–45	Cross-sectional, comparative	SAC	Eyetracker	Cross	No	70 cm
2018	Rizzo ⁴²	42 normal subjects aged 19–52	Cross-sectional, comparative	SAC	Eyetracker and K-D	Numbers on test card	No	Not described
2018	Wetzel ⁴³	75 normal subjects aged 21–53	Cross-sectional, comparative	SAC	Eyetracker	Nine separated points	No	75 cm
2019	Hoffmann ⁴⁴	40 normal subjects	Cross-sectional, comparative	SAC	Eyetracker	Circle	No	Not described
2019	Chehrehnegar ⁴⁵	59 normal subjects	Cross-sectional, comparative	SAC	Eyetracker	Cross (Simon task)	No	60 cm
2019	Wetzel ⁴⁶	75 normal subjects aged 18–65	Cross-sectional, comparative	SAC, SP	Eyetracker	Nine separated points	No	75 cm
2020	Wertli ⁴⁷	25 normal subjects aged 10–11	Cross-sectional, descriptive	SAC, FIX	Eyetracker	Text	Yes	Between 50 to 80 cm
2020	Sinno ¹⁹	120 normal subjects aged 5–17	Cross-sectional, descriptive	SAC, SP	Videonystagmography	White Square	No	120 cm
2021	Hindmarsh ⁴⁸	196 subjects aged 7.3–8.9	Cross-sectional, descriptive	SAC, SP, FIX	Eyetracker and DEM	Numbers on test card	Yes	60 cm
2021	Kullmann ⁴⁹	466 normal subjects aged 18–45	Cross-sectional, descriptive	SAC, SP	Eyetracker	Dot	No	Not described
2021	Amato ⁵⁰	54 normal subjects aged 18–28	Pseudoexperimental	SAC	Eyetracker	Videogame	No	Not described
2022	D'Addio ⁵¹	34 normal subjects	Cross-sectional, comparative	FIX	Eyetracker	White disc in screen	No	65 cm
2022	Eichler ⁵²	52 normal subjects aged 9–34	Cross-sectional, descriptive	SAC	Eyetracker	OMAT	Yes	25 cm
2023	Orduna-Hospital ⁵³	52 normal subjects aged 18–30	Cross-sectional, descriptive	SAC, FIX	Eyetracker and DEM	Numbers on test card	Yes	60 cm

Table 1 (Continued)	Year	Author	Characteristics of the control group (n, ranged age)	Study design	Oculomotor skills	Measurments system	Stimulus	Optometric examination	Work Distance (cm)
	2023	Wertli ⁵⁴ SAC: saccadic SP: smooth pursuits FIX: fixation	118 normal subjects aged 7–12	Cross-sectional, descriptive	SAC, FIX DEM: developmental eye movement OMAT: oculomotor assessment tool test	Eyetracker	Text	Yes	Between 50 to 80 cm

Table 2 Most commonly measured aspects of oculomotor parameters in the articles reviewed.

Skills	Aspects	n	%
Saccades	Number of saccades	11	28.9
	Amplitude of saccades	8	21.1
	Velocity of saccades	8	21.1
	Latency of saccades	11	28.9
Smooth pursuits	Pursuit gain	10	26.3
Fixation	Fixation time	9	23.7

studies (5.3 %) provided a range but did not specify an exact value.

Oculomotor skills aspects

Table 2 presents the aspects of oculomotor skills measured for each parameter (saccadic, smooth pursuit, and fixation movements), extracted from the studies included in this systematic review. It is evident that there is considerable diversity in the parameters analyzed for each test. Additionally, it can be observed that most of these parameters are not analyzed in >20 % of the articles.

For saccadic movements, the most frequently analyzed parameters are the *number of saccades* (28.9 %), *latency* (28.9 %), *velocity* (21.1 %) and *amplitude* (21.1 %). In pursuits movements, the most commonly measured aspect is *pursuit gain* (26.3 %). Lastly, for fixation movements, the most frequently measured parameter is *fixation time* (23.7 %). In addition, authors did not find reported data on Bivariate Contour Ellipse Area (BCEA), a standard metric of fixation stability.

Other aspects were also measured by the authors of the included articles, such as *Peak Velocity* in saccadic movements, *velocity* in smooth pursuits and *number of fixations* in fixation movements. However, these aspects were measured less frequently, in fewer than 20 % of the articles.

Assessment of quality and bias of included studies

Table 3 and Fig. 2 present the analysis of the 38 included studies using the QUADAS-2 tool, which evaluates the risk of bias and applicability. In general, it can be observed that there is a high risk of bias (≥ 50 % in the aspects of “flow and timing”, “reference standard” and “patient selection,” as well as poor applicability (≥ 50 %), across all aspects.

Discussion

Oculomotor function has gained increasing relevance in recent years. In fact, more than half of the studies analyzed in this review (20 out of 38; 52.6 %) have been conducted in the last decade.^{19,36–54}

Firstly, the technology used to measure oculomotor function has advanced considerably. Early studies relied on

Table 3 Quality rating of the 38 included studies in the systematic review (QUADAS-2 results) (☺: low risk; ☹: high risk; ?: unclear risk.).

Study	Risk of bias				Applicability concerns		
	Patient selection	Index test	Reference standard	Flow and timing	Patient selection	Index test	Reference standard
1980 Schalen ²¹	☹	☹	?	☺	☹	☹	☺
1984 Berginus ²²	☺	☺	?	?	☹	☹	☺
1990 Garzia ¹⁴	☹	☺	?	☺	☹	☹	☺
1992 Maples ¹⁵	☹	☹	?	☺	☹	☹	☺
1992 Versino ²³	☹	☺	?	☹	☹	☹	☺
1993 Ross ²⁴	☹	☺	?	☹	☹	☺	☺
1997 Litman ²⁵	☹	☹	?	☹	☹	?	☹
1997 Ross ²⁶	☹	?	☹	☹	☹	☺	☹
1999 Campana ²⁷	☹	☺	☹	☹	☹	☹	☹
2003 Jimenez ²⁸	☺	?	☺	☹	☺	☺	☺
2006 Rütsche ¹⁸	☺	☺	☹	☹	☺	☹	☹
2007 Aring ²⁹	☺	☺	☹	☹	☺	☺	☺
2010 van Tritch ³⁰	☹	☹	☹	☹	☹	☹	☹
2011 Goepel ³¹	☹	?	☹	☹	☹	☹	☹
2011 Kattoulas ³²	?	☺	☹	☹	?	☺	☺
2011 Webber ³³	☺	☺	☺	☹	☺	☺	☺
2012 Shi ³⁴	?	☺	☹	?	☺	?	☹
2012 Boot ³⁵	☺	☺	☹	☹	☺	?	☹
2015 Seferlis ³⁶	☺	☺	☹	☹	☺	☺	☺
2015 Doetti ³⁷	☺	☺	☹	☹	?	?	☹
2016 Zalla ³⁸	☹	☺	☹	☹	?	☹	☹
2016 Yang ³⁹	☹	☺	☹	☹	☹	?	☺
2017 Choi ⁴⁰	☹	☹	☹	☹	☹	☹	☹
2017 Ferreira ⁴¹	☹	☹	☹	☹	☹	☹	☹
2018 Rizzo ⁴²	?	☺	?	☹	☹	☹	☹
2018 Wetzel ⁴³	☹	☺	☹	☹	☹	☹	☹
2019 Hoffman ⁴⁴	☹	☺	☹	☹	☹	☹	☹
2019 Chehrehnegar ⁴⁵	☹	☺	☹	☹	☹	☹	☹
2019 Wetzel ⁴⁶	☹	☺	☹	☹	?	☹	☹
2020 Wertli ⁴⁷	☺	☺	☹	☹	?	?	☺
2020 Sinno ¹⁹	☺	☺	☹	☹	☺	☺	☺
2021 Hindmarsh ⁴⁸	☺	☺	☹	☹	?	☺	☺
2021 Kullman ⁴⁹	☺	☺	☹	☹	?	☺	☺
2021 Amato ⁵⁰	☹	☹	☹	☹	☹	☹	☹
2022 D'Addio ⁵¹	☹	☹	☹	☹	☹	☹	☹
2022 Eichler ⁵²	☺	☺	☹	☹	☺	☺	☺
2023 Orduna-Hospital ⁵³	☺	☺	☺	☹	☺	?	☺
2023 Wertli ⁵⁴	☺	☺	☹	☹	?	?	☺




Fig. 2 QUADAS-2 domain for articles included in the systematic review.

electrooculography and subjective assessments, whereas more recent research utilizes eye-tracking systems and video-nystagmography (Table 1). This shift suggests a notable improvement in the quantitative analysis of ocular motor skills. Secondly, it is now demonstrated that oculomotor disturbances may be present in some systemic conditions, such as neurological diseases^{3-6,50} or development disorders.⁷ Despite current devices seeming to measure oculomotor function in a better way, it is not completely demonstrated because there is a great methodological variability between studies in terms of the type of stimulus, the complexity of the task, the working distance, and even the analyzed parameters. Such variability precludes direct comparison of measurements across different systems and constrains the ability to perform agreement analyses between devices. Consequently, there are very few studies comparing subjective methods (DEM, K-D, NSUCO) with objective methods based on eyetrackers,^{33,42,48} and even fewer comparing several objective methods with one another. Not only are comparisons between devices not adequately developed in scientific literature, but also the validation with a single device is also insufficient because typically only one measurement is taken for each subject, preventing the development of repeatability analysis. In fact, only the DEM¹⁴ and NSUCO¹⁵ techniques have been thoroughly validated by measuring twice for each subject and calculating intrasession repeatability. The impossibility of generalizing the results is also supported by the study design and population. Oculomotor function is highly age-dependent,^{14,15,18,19,32,36,55} and there are multiple studies involving adults, children or both. In contrast, almost half of the studies analyzed in this paper (15 of 38 studies; 39.5 %)^{25-27,30,34,37,38,40-46,51} are cross-sectional with comparative design incorporating a control group whose ocular movements are compared to those obtained by a specific group with a particular condition or disorder. This involves selecting the control group based on the study group's characteristics, inducing a patient selection bias that prevents the results from being generalized to the broader population. Another finding observed in this systematic review, potentially affecting the inference process, is the lack of an optometric evaluation in the majority of studies (26 of 38 studies; 68.4 %).^{14,15,19,21-27,30-32,35,}

^{38-46,49-51} Among 12 articles that conducted previous eye tests,^{18,28,29,33,34,36,37,47,48,52-54} only one study performed a comprehensive optometric examination evaluating accommodation, vergences, and stereopsis.⁵³ Eight articles conducted some accommodative and binocular tests, but these were insufficient for the purposes of this study,^{18,28,29,33,47,48,52,54} and the other three articles only included VA measures.^{34,36,37} It represents a very remarkable limitation of the current evidence because an evaluation of refraction, accommodation and vergences would be elementary in this topic as indicated Cacho-Martínez et al. (2024 Cacho-Martínez) Moreover, difficulty in focusing or maintaining single (haplopic) vision is related to the efficiency of fixations and saccades. Therefore, a prior optometric examination is important to detect binocular or accommodative dysfunctions that may affect oculomotor performance (REF 2022, Liu Z)

It is reported that oculomotor function is impaired in individuals with amblyopia⁵⁶ and strabismus,⁵⁷ affecting both the dominant and amblyopic eye.⁵⁸ Additionally, monocular and binocular oculomotor skills differ.⁵⁹ In cases of convergence insufficiency, alterations in reading saccades are also observed.⁶⁰ Difficulties in near vision due to accommodative issues and/or uncorrected refractive errors may also contribute to oculomotor dysfunction, as suggested by Liu et al.⁶¹ Therefore, a comprehensive visual examination that includes binocular, accommodative, and sensory assessments is necessary to rule out the presence of amblyopia, strabismus, non-strabismic binocular dysfunctions, and oculomotor dysfunctions. Such an approach would reduce the number of visual variables that interfere with measurement and facilitate the establishment of normative oculomotor parameters. It is also essential to measure monocular and binocular visual acuity at both distance and near. Furthermore, it should be noted that the presence of strabismus can result in a number of abnormalities across numerous brain areas involved in visual functions and eye movements.^{62,63} Indeed, not all eye tracker software is designed for the precise measurement of strabismus. Some software merely detects the initial position of the eyes during calibration, without the capability to distinguish whether the eyes are aligned or if strabismus is present. Considering

that oculomotor function is altered when patients do not use the best optical correction or have other visual anomalies,⁶⁴ the low rate of studies including an optometric evaluation in their sample represents an important limitation (2024 Cacho-Martínez). Other potential biases and applicability concerns have been analyzed in this paper using the QUADAS-2 tool (Table 3, Fig. 2).

Regarding the methods used to measure oculomotor function in the articles included in this systematic review, Table 1 highlights the presence of various stimuli, measurement distances, technologies, and types of visual tasks. Starting with the stimuli employed, a wide range of tests was used, including classic tests such as DEM, NSUCO, K-D, or reading a text, as well as the use of computers to project points, shapes, letters, or cartoons, and even a video game or the OculoMotor Assessment Tool (OMAT). This variety of stimuli, differing in shape, size, color, and cognitive processing difficulty, represents a limitation when attempting to compare them. In fact, the cognitive processes required to follow a moving point differ significantly from those involved in reading text, numbers, or shapes, and are distinct from observing drawings or playing a video game.¹² Therefore, this variability constitutes a critical factor when measuring oculomotor function.

During the measurement process, attention loss due to the appearance of another stimulus or, in the case of computers or screens, the presence of additional objects can influence the results. Similarly, the working distance is another limitation. As shown in Table 1, four studies (10.5 %) performed measurements at a near distance (≤ 40 cm),^{14,18,28,52} 18 studies (47.4 %) at an intermediate distance (between 40 and 100 cm),^{22,24,25,27,29,31,35,38,40,41,43,45–48,51,53,54} seven studies (18.4 %) at a medium-far distance (≥ 100 cm),^{19,21,24,34,36,37,39} and nine studies (23.7 %) did not specify the working distance.^{15,23,30,32,33,42,44,49,50} Although no gold standard exists for the working distance, it can affect the accuracy of eye movement recordings. While it is true that the eye's angular orientation relative to the stimulus is key—and at different distances but the same angular orientation, the eyes should theoretically behave similarly in their movement—factors such as peripheral vision capacity or the limitation of specific ocular muscles may affect the measurement. At greater distances, these factors could introduce errors. These errors are due to the interaction between the accommodative and vergence system,^{65,66} and it has been shown in scientific literature that variation in working distance produces variation in the accommodative and vergence system.⁶⁷

Regarding the technology used, there has been significant progress from the early methods, such as electrooculography (an invasive technique), to the development of video nystagmography, video oculography, and the most widely used technology: eye trackers. The use of eye trackers in studies has grown exponentially, particularly in the optometric field, enabling more precise and objective measurements of oculomotor function. However, it is essential to consider that each technology differs in its stimuli, working distances, and limitations, complicating comparisons between studies due to the lack of a gold standard for measurement.

Other factors that may influence results include the ergonomics of the measurement systems. Modern systems, such as eye trackers or video-oculographs, exhibit considerable

variability, including trial frames with side-mounted cameras, helmets, virtual reality glasses, and bars placed on computer screens. Additionally, some devices are not commercially available but are laboratory prototypes or optical bench setups. Establishing unified ergonomic criteria would be beneficial.

During the evaluation of oculomotor function, it is essential to consider the subject's age. Scientific literature has demonstrated that oculomotor function varies with age, and studies that propose normative values present results based on age.^{19,47–49,52–54} Research suggests that oculomotor performance improves during the early years of life (approximately from 4 to 15 years)^{68–70} and subsequently declines.^{55,70,71} Another critical aspect to consider is the subject's cognitive level. Neurodevelopmental disorders,^{72,73} neurological diseases,^{74,75} or, as mentioned earlier, age—particularly in early childhood and older adulthood—can significantly influence the measurements.

Furthermore, maintaining the subject's attention is crucial. Oculomotor measurements typically last between 30 s and 5 min, and any loss of attention during the evaluation can lead to inconsistent or unreliable results. Similarly, fatigue and exhaustion during the test—especially in lengthy, tedious evaluations involving complex stimuli or requiring high cognitive demand—can introduce biases into the measurements.⁷⁶ A clear parallel can be observed in visual field testing, where efforts have been made for years to shorten the test duration to mitigate these biases.

All these factors—age, cognitive level, attention, fatigue, and exhaustion—must be carefully considered when establishing a gold standard for oculomotor function measurement. Unified criteria should be defined to ensure consistency and reliability in the evaluation process.

Following the identification of methodological considerations frequently neglected in the literature and warranting careful attention, it is equally imperative to undertake a critical appraisal of the overall quality and methodological rigor of the included studies. In this systematic review, this analysis was conducted using the QUADAS-2 tool (Table 3, Fig. 2), which demonstrated a high incidence of studies with significant risks of bias and important limitations regarding applicability concerns. In particular, the most common risk of bias is related to reference standards, as only 4 out of 38 studies (10.5 %)^{28,33,48,53} compare the test under study with a gold standard technique or at least other independent method capable of measuring similar parameters of oculomotor function. This lack of comparison prevents validation of the measurements due to the impossibility of performing an agreement analysis. Similarly, flow and timing are also highly affected by a high risk of bias in almost all studies, with only 3 out of 38 studies (7.9 %)^{14,15,21} being free of this type of bias. The bias was generally caused by the predominance of single measurements per device, which prevents repeatability analysis. This finding further contributes to the generation of non-validated measurements that cannot be used as normative values.

Patient selection is another significant source of bias present in most studies, as many included subjects with highly specific characteristics—for example, only highly educated individuals⁴⁰ or only students⁵⁰—limiting extrapolation to the general population. Additionally, the selection of

normal subjects was often influenced by the demographics and characteristics of other study groups involved in comparative studies. Therefore, recruitment in these comparative studies is not randomized. Furthermore, comparative studies often attempt to match ages between groups, leading to significant heterogeneity among control groups depending on the nature of the study.

On the other hand, it is important to highlight the higher-quality results regarding measurement methodology or index test in the current review, as 27 out of 38 studies (71.1 %) provide a complete description of the procedure.^{14,18,19,22–24,27,29,32–39,42–49,52–54} However, despite an accurate description of the measurement process, technology, and tasks, in some cases, the methodology remains difficult to replicate, affecting its applicability. This limitation is primary due to the use of outdated technology,^{21–23} implementation of subjective tests,¹⁵ non-standardized texts that introduce cognitive and educational dependencies,^{40,54} and highly complex tests that are difficult to replicate, such as the Convirt Test⁵⁰ or OMAT Test.⁵²

Applicability concerns are further compounded by unclear or questionable inclusion criteria for subjects, as well as the lack of proper characterization of measurements, leading to data that do not contribute to establishing normative values.

Regarding the limitations of the present study, the scarcity of specific articles evaluating normative values must be noted, as most studies focus on comparative or interventional approaches rather than the characterization of oculomotor function. Additionally, intrinsic limitations of the current tools used to assess study quality may lead to incomplete considerations or misinterpretations, even though QUADAS-2 remains the most appropriate tool for the objectives of this investigation. Because of these limitations a unique standardization of oculomotor measurements is currently not possible. Future studies should be conducted to propose a preliminary oculomotor protocol by an expert committee which should be validated in different sample characteristics.

On the other hand, key strengths of this study include its innovative and applicable focus, emphasizing the need to establish normative values. Moreover, the development and analysis of the procedure were conducted independently by three investigators, with a final consensus to enhance the evaluation of the current evidence.

Conclusion

Based on these findings, this systematic review concludes that there is no clear evidence on normative values for oculomotor skills, nor is there a consensus on the measurement methods, stimuli used, or working distances. Furthermore, there is no agreement on which specific aspects of oculomotor skills should be assessed, and due to the quality evaluation of these studies, the risk of bias remains predominantly high.

Additionally, it is concerning that the vast majority of studies do not include a comprehensive optometric assessment – incorporating refractive, accommodative, and binocular tests – which may introduce bias in the results. To enhance reliability and applicability, measurement criteria

should be standardized, and normative values should be established.

Declaration of competing interest

The authors have no conflicts of interest to declare.

Funding

No funding.

Supplementary materials

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.optom.2025.100570](https://doi.org/10.1016/j.optom.2025.100570).

References

- Doettl SM, McCaslin DL. Oculomotor assessment in children. *Semin Hear*. 2018;39:275–287.
- Salgado-Fernandez A, Vazquez-Amor A, Alvarez-Peregrin C, Martinez-Perez C, Villa-Collar C, Angel Sanchez-Tena M. Influence of eye movements on academic performance: a bibliometric and citation network analysis. *J Eye Mov Res*. 2022;15.
- Wolf A, Ueda K, Hirano Y. Recent updates of eye movement abnormalities in patients with schizophrenia: a scoping review. *Psychiatry Clin Neurosci*. 2021;75:82–100.
- Wolf A, Tripanpitak K, Umeda S, Otake-Matsuura M. Eye-tracking paradigms for the assessment of mild cognitive impairment: a systematic review. *Front Psychol*. 2023;14:1197567.
- Noyes B, Biorac A, Vazquez G, Khalid-Khan S, Munoz D, Booi L. Eye-tracking in adult depression: protocol for a systematic review and meta-analysis. *BMJ Open*. 2023;13:e069256.
- Sekar A, Panouilleres MTN, Kaski D. Detecting abnormal eye movements in patients with neurodegenerative diseases: current insights. *Eye Brain*. 2024;16:3–16.
- Bilbao C, Pinero DP. Clinical characterization of oculomotricity in children with and without specific learning disorders. *Brain Sci*. 2020;10.
- Vinuela-Navarro V, Goset J, Aldaba M, et al. Eye movements in patients with post-COVID condition. *Biomed Opt Express*. 2023;14:3936–3949.
- Gheorghe CM, Purcarea VL, Gheorghe IR. Using eye-tracking technology in Neuromarketing. *Rom J Ophthalmol*. 2023;67:2–6.
- Spataro R, Ciriaco M, Manno C, La Bella V. The eye-tracking computer device for communication in amyotrophic lateral sclerosis. *Acta Neurol Scand*. 2014;130:40–45.
- Gonzalez-Vides L, Hernandez-Verdejo JL, Canadas-Suarez P. Eye tracking in optometry: a systematic review. *J Eye Mov Res*. 2023;16. <https://doi.org/10.16910/jemr.16.3.3>.
- Nieboer W, Ghiani A, de Vries R, Brenner E, Mann DL. Eye tracking to assess the functional consequences of vision impairment: a systematic review. *Optom Vis Sci*. 2023;100:861–875.
- Pinero DP. Oculomotor dysfunctions: evidence-based practice. *J Optom*. 2020;13:137–138.
- Garzia RP, Richman JE, Nicholson SB, Gaines CS. A new visual-verbal saccade test: the development eye movement test (DEM). *J Am Optom Assoc*. 1990;61:124–135.

15. Maples WC, Atchley J, Ficklin T. Northeastern State University College of Optometry's oculomotor norms. *J Behav Optom*. 1992;3:143–150.
16. Lieberman S, Cohen AH, Rubin J. NYSOA K-D test. *J Am Optom Assoc*. 1983;54:631–637.
17. Curthoys IS, Ooi S, Wearne SL, Todd MJ, Halmagyi GM. Normative data for human smooth pursuit responses to horizontal step-ramp targets. *Acta Otolaryngol Suppl*. 1991;481:391–393.
18. Rutsche A, Baumann A, Jiang X, Mojon DS. Development of visual pursuit in the first 6 years of life. *Graefes Arch Clin Exp Ophthalmol*. 2006;244:1406–1411.
19. Sinno S, Najem F, Abouchakra KS, Perrin P, Dumas G. Normative values of saccades and smooth pursuit in children aged 5 to 17 years. *J Am Acad Audiol*. 2020;31:384–392.
20. Wilson SJ, Glue P, Ball D, Nutt DJ. Saccadic eye movement parameters in normal subjects. *Electroencephalogr Clin Neurophysiol*. 1993;86:69–74.
21. Schalen L. Quantification of tracking eye movements in normal subjects. *Acta Otolaryngol*. 1980;90:404–413.
22. Bergenius J. Computerized analysis of voluntary eye movements. A clinical method for evaluation of smooth pursuit and saccades in oto-neurological diagnosis. *Acta Otolaryngol*. 1984;98:490–500.
23. Versino M, Grassi M, Genovese E, Zambarbieri D, Schmid R, Cosi V. Quantitative evaluation of saccadic eye movements. *Neuro-Ophthalmol*. 1992;12:327–342.
24. Ross RG, Radant AD, Hommer DW. A developmental study of smooth pursuit eye movements in normal children from 7 to 15 years of age. *J Am Acad Child Adolesc Psychiatry*. 1993;32:783–791.
25. Litman RE, Torrey EF, Hommer DW, Radant AR, Pickar D, Weinberger DR. A quantitative analysis of smooth pursuit eye tracking in monozygotic twins discordant for schizophrenia. *Arch Gen Psychiatry*. 1997;54:417–426.
26. Ross DE, Thaker GK, Buchanan RW, et al. Eye tracking disorder in schizophrenia is characterized by specific ocular motor defects and is associated with the deficit syndrome. *Biol Psychiatry*. 1997;42:781–796.
27. Campana A, Duci A, Gambini O, Scarone S. An artificial neural network that uses eye-tracking performance to identify patients with schizophrenia. *Schizophr Bull*. 1999;25:789–799.
28. Jimenez R, Gonzalez MD, Perez MA, Garcia JA. Evolution of accommodative function and development of ocular movements in children. *Ophthalmic Physiol Opt*. 2003;23:97–107.
29. Aring E, Gronlund MA, Hellstrom A, Ygge J. Visual fixation development in children. *Graefes Arch Clin Exp Ophthalmol*. 2007;245:1659–1665.
30. van Tricht MJ, Nieman DH, Bour LJ, et al. Increased saccadic rate during smooth pursuit eye movements in patients at ultra high risk for developing a psychosis. *Brain Cogn*. 2010;73:215–221.
31. Goepel J, Biehl SC, Kissler J, Paul-Jordanov I. Pro- and antisaccades in children elicited by visual and acoustic targets - does modality matter? *BMC Pediatr*. 2011;11:116.
32. Kattoulas E, Smyrnis N, Stefanis NC, Avramopoulos D, Stefanis CN, Evdokimidis I. Predictive smooth eye pursuit in a population of young men: I. Effects of age, IQ, oculomotor and cognitive tasks. *Exp Brain Res*. 2011;215:207–218.
33. Webber A, Wood J, Gole G, Brown B. DEM test, visagraph eye movement recordings, and reading ability in children. *Optom Vis Sci*. 2011;88:295–302.
34. Shi XF, Xu LM, Li Y, Wang T, Zhao KX, Sabel BA. Fixational saccadic eye movements are altered in anisometropic amblyopia. *Restor Neurol Neurosci*. 2012;30:445–462.
35. Boot FH, Pel JJ, Evenhuis HM, van der Steen J. Quantification of visual orienting responses to coherent form and motion in typically developing children aged 0-12 years. *Invest Ophthalmol Vis Sci*. 2012;53:2708–2714.
36. Seferlis F, Chimona TS, Papadakis CE, Bizakis J, Triaridis S, Skoulakis C. Age related changes in ocular motor testing in healthy subjects. *J Vestib Res*. 2015;25:57–66.
37. Doettl SM, Plyler PN, McCaslin DL, Schay NL. Pediatric oculomotor findings during monocular videonystagmography: a developmental study. *J Am Acad Audiol*. 2015;26:703–715.
38. Zalla T, Seassau M, Cazalis F, Gras D, Leboyer M. Saccadic eye movements in adults with high-functioning autism spectrum disorder. *Autism*. 2018;22:195–204.
39. Yang CJ, Lee JY, Kang BC, Lee HS, Yoo MH, Park HJ. Quantitative analysis of gains and catch-up saccades of video-head-impulse testing by age in normal subjects. *Clin Otolaryngol*. 2016;41:532–538.
40. Choi W, Lowder MW, Ferreira F, Swaab TY, Henderson JM. Effects of word predictability and preview lexicality on eye movements during reading: a comparison between young and older adults. *Psychol Aging*. 2017;32:232–242.
41. Ferreira M, Pereira PA, Parreira M, et al. Using endogenous saccades to characterize fatigue in multiple sclerosis. *Mult Scler Relat Disord*. 2017;14:16–22.
42. Rizzo JR, Hudson TE, Amorapant PX, et al. The effect of linguistic background on rapid number naming: implications for native versus non-native english speakers on sideline-focused concussion assessments. *Brain Inj*. 2018;32:1690–1699.
43. Wetzel PA, Lindblad AS, Raizada H, et al. Eye tracking results in postconcussive syndrome versus normative participants. *Invest Ophthalmol Vis Sci*. 2018;59:4011–4019.
44. Hoffmann A, Ettinger U, Montoro C, Reyes Del Paso GA, Duschek S. Cerebral blood flow responses during prosaccade and antisaccade preparation in major depression. *Eur Arch Psychiatry Clin Neurosci*. 2019;269:813–822.
45. Chehrehnegar N, Nejati V, Shati M, et al. Behavioral and cognitive markers of mild cognitive impairment: diagnostic value of saccadic eye movements and Simon task. *Aging Clin Exp Res*. 2019;31:1591–1600.
46. Wetzel PA, Lindblad AS, Mulatya C, et al. Eye tracker outcomes in a randomized trial of 40 sessions of hyperbaric oxygen or sham in participants with persistent post concussive symptoms. *Undersea Hyperb Med*. 2019;46:299–311.
47. Wertli J, Schotzau A, Trauzettel-Klosinski S, Palmowski-Wolfe A. Feasibility of eye movement recordings with the SMI tracking bar in 10- to 11-year-old children performing a reading task. *Klin Monbl Augenheilkd*. 2020;237:510–516.
48. Hindmarsh GP, Black AA, White SL, Hopkins S, Wood JM. Eye movement patterns and reading ability in children. *Ophthalmic Physiol Opt*. 2021;41:1134–1143.
49. Kullmann A, Ashmore RC, Braverman A, et al. Portable eye-tracking as a reliable assessment of oculomotor, cognitive and reaction time function: normative data for 18-45 year old. *PLoS One*. 2021;16:e0260351.
50. Amato I, Nanav A, Piantella S, et al. Assessing the utility of a virtual-reality neuropsychological test battery, 'CONVIRT', in detecting alcohol-induced cognitive impairment. *Behav Res Methods*. 2021;53:1115–1123.
51. D'Addio F, Pastore I, Loretelli C, et al. Abnormalities of the oculomotor function in type 1 diabetes and diabetic neuropathy. *Acta Diabetol*. 2022;59:1157–1167.
52. Eichler R, Mivtachi G, Hershkovitz-Azoulay D, Scheiman M, Ben-Eli H. OculoMotor assessment tool: children compared with adults. *Clin Optom (Auckl)*. 2022;14:75–81.
53. Orduna-Hospital E, Navarro-Marques A, Lopez-de-la-Fuente C, Sanchez-Cano A. Eye-tracker study of the developmental Eye movement test in young people without binocular dysfunctions. *Life (Basel)*. 2023;13(3):773.
54. Wertli J, Schotzau A, Palmowski-Wolfe A. The influence of age on eye movements during reading in early elementary school children. *Klin Monbl Augenheilkd*. 2023;240:591–598.

55. Pueyo V, Yam JCS, Perez-Roche T, et al. Development of oculomotor control throughout childhood: a multicenter and multi-ethnic study. *J Vis.* 2022;22:4.
56. Webber AL. The functional impact of amblyopia. *Clin Exp Optom.* 2018;101:443–450.
57. Hepokur M, Mutlu B, Gunes M, et al. The effect of different types of convergent strabismus on horizontal eye movements. *Int Ophthalmol.* 2022;42:3951–3961.
58. Birch EE, Kelly KR, Giaschi DE. Fellow eye deficits in amblyopia. *J Binocul Vis Ocul Motil.* 2019;69:116–125.
59. Otero-Millan J, Macknik SL, Martinez-Conde S. Fixational eye movements and binocular vision. *Front Integr Neurosci.* 2014;8:52.
60. Hirota M, Kanda H, Endo T, et al. Relationship between reading performance and saccadic disconjugacy in patients with convergence insufficiency type intermittent exotropia. *Jpn J Ophthalmol.* 2016;60:326–332.
61. Liu Z, Zhang K, Gao S, Yang J, Qiu W. Correlation between eye movements and asthenopia: a prospective observational study. *J Clin Med.* 2022;11(23):7043.
62. Metz HS. Restrictive factors in strabismus. *Surv Ophthalmol.* 1983;28:71–83.
63. Walton MMG, Pallus A, Fleuriet J, Mustari MJ, Tarczy-Hornoch K. Neural mechanisms of oculomotor abnormalities in the infantile strabismus syndrome. *J Neurophysiol.* 2017;118:280–299.
64. Scheiman M, Wick B. *Clinical Management of Binocular Vision. Heterophoric, Accommodative, and Eye Movement Disorders.* Philadelphia: Wolters Kluwer; 2019.
65. Hung GK. Adaptation model of accommodation and vergence. *Ophthalmic Physiol Opt.* 1992;12:319–326.
66. Hung GK, Ciuffreda KJ, Rosenfield M. Proximal contribution to a linear static model of accommodation and vergence. *Ophthalmic Physiol Opt.* 1996;16:31–41.
67. Rosenfield M. Tonic vergence and vergence adaptation. *Optom Vis Sci.* 1997;74:303–328.
68. Accardo AP, Pensiero S, Da Pozzo S, Perissutti P. Characteristics of horizontal smooth pursuit eye movements to sinusoidal stimulation in children of primary school age. *Vision Res.* 1995;35:539–548.
69. Purves D, Augustine GJ, Fitzpatrick D, et al. Types of eye movements and their functions. *Neuroscience.* Sunderland, MA, USA: Sinauer Associates; 2001.
70. Luna B, Velanova K, Geier CF. Development of eye-movement control. *Brain Cogn.* 2008;68:293–308.
71. Blignaut P, van Rensburg EJ, Oberholzer M. Visualization and quantification of eye tracking data for the evaluation of oculomotor function. *Heliyon.* 2019;5:e01127.
72. Bilbao C, Pinero DP. Objective and subjective evaluation of saccadic eye movements in healthy children and children with neurodevelopmental disorders: a pilot study. *Vision (Basel).* 2021;5(2):28.
73. Bilbao C, Pinero DP. Distribution of visual and oculomotor alterations in a clinical population of children with and without neurodevelopmental disorders. *Brain Sci.* 2021;11(3):351.
74. Caldani S, Amado I, Bendjemaa N, et al. Oculomotricity and neurological soft signs: can we refine the endophenotype? A study in subjects belonging to the spectrum of schizophrenia. *Psychiatry Res.* 2017;256:490–497.
75. Gil-Casas A, Pinero DP, Molina-Martin A. Are near visual signs and symptoms in multiple sclerosis compatible with convergence insufficiency? *Clin Exp Optom.* 2022;105:631–636.
76. Bruce BB, Newman NJ. Functional visual loss. *Neurol Clin.* 2010;28:789–802.