



ORIGINAL ARTICLE

Positive relative accommodation is an independent risk factor for myopia onset: a prospective cohort study among chinese primary schoolchildren, the WEPrOM study



Yingying Ye^{a,b,c,1}, Yimin Yuan^{a,b,c,1}, Chengnan Guo^{e,1}, Yingying Huang^{a,b,c},
Jingwei Zheng^{a,b}, Yee Ling Wong^{c,d}, Binbin Su^{a,b,c}, Yang Ding^{a,b,c}, Björn Drobe^{c,d},
Minfeng Chen^a, Hao Chen^{a,b,1,*}, Jinhua Bao^{a,b,c,1,*}

^a National Engineering Research Centre of Ophthalmology and Optometry, Eye Hospital, Wenzhou Medical University, Wenzhou, Zhejiang, China

^b National Clinical Research Center for Ocular Diseases, Wenzhou, Zhejiang, China

^c WEIRC, Wenzhou Medical University-Essilor International Research Centre, Wenzhou Medical University, Wenzhou, Zhejiang, China

^d Research and Development, EssilorLuxottica, Singapore, Singapore

^e Wenzhou Medical University, Wenzhou, Zhejiang, China

Received 7 April 2025; accepted 25 July 2025

Available online 30 August 2025

KEYWORDS

Myopia onset;
Risk factors;
Schoolchildren;
Cohort study;
Positive relative accommodation

Abstract

Purpose: To identify independent risk factors for myopia onset in schoolchildren, with a focus on binocular visual function.

Methods: We conducted a school-based prospective cohort study in Wenzhou, China. Schoolchildren in grades 2 and 3 were recruited in 2014 and followed until graduation at grade 6. Myopia was defined as a spherical equivalent refraction (SER) of ≤ -0.50 diopters. The risk factors assessed included monocular uncorrected visual acuity (UCVA), axial length (AL), corneal refractive power (CR), demographic characteristics, daily activities, parental myopia, parental education level, and routine clinical binocular visual function parameters such as phoria, accommodation, and convergence-related metrics.

Results: Multivariable logistic regression analysis revealed that children with the following baseline characteristics had a significantly increased risk (OR;95% CI) of developing myopia before

List of abbreviations: WEPrOM, Wenzhou Medical University Essilor Progression and Onset of Myopia; UCVA, uncorrected visual acuity; log-MAR, logarithm of the minimum angle of resolution; SER, spherical equivalent refraction; D, diopter; AL, axial length; CR, corneal refractive power; AC/A, accommodative convergence to accommodation; NRA, negative relative accommodation; PRA, positive relative accommodation; BI, base-in; BO, base-out; PD, prism diopter.

* Correspondence authors at: Eye Hospital, School of Ophthalmology and Optometry, Wenzhou Medical University, 270 Xueyuan Road, Wenzhou, Zhejiang, 325027, China.

E-mail addresses: chenhao@mail.eye.ac.cn (H. Chen), baojessie@mail.eye.ac.cn (J. Bao).

¹ These authors contributed equally to this work.

<https://doi.org/10.1016/j.optom.2025.100577>

1888-4296/© 2025 The Authors. Published by Elsevier España, S.L.U. on behalf of Spanish General Council of Optometry. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

graduation: female sex (3.03;1.99–4.62; $P<.001$), having two myopic parents (2.36;1.29–4.31; $P=.005$), worse UCVA (19.99;2.24–178.44; $P=.007$), more negative SER values (0.15;0.07–0.31; $P<.001$), longer AL (7.28;4.30–12.31; $P<.001$), larger CR (2.20;1.75–2.76; $P<.001$), and lower magnitude of positive relative accommodation (PRA) (1.11;1.02–1.22; $P=.02$). Additional exploratory subgroup analyses indicated that the association between PRA and myopia incident remained consistent across various demographic characteristics (P -interaction >0.05). Receiver operating characteristic curves (AUC; 95% CI) demonstrated that PRA (0.59;0.55–0.63) exhibited predictive capability comparable to key ocular biometric parameters such as AL (0.57;0.53–0.62) and CR (0.58;0.53–0.62).

Conclusions: The current study identifies PRA as a stable, independent risk factor for myopia onset, with predictive capability comparable to key ocular biometric parameters. This finding can be utilized in future studies to enhance the accuracy of myopia prediction and assist in making informed decisions regarding myopia interventions.

© 2025 The Authors. Published by Elsevier España, S.L.U. on behalf of Spanish General Council of Optometry. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

In recent decades, there has been a significant global increase in the severity of myopia among schoolchildren, particularly in East and Southeast Asia, where the condition has reached epidemic proportions.^{1–3} In China, the annual incidence of myopia among schoolchildren has been reported to range from 6.3% to 31.7%.^{4–11} Myopia and high myopia can lead to permanent pathological changes and result in both direct healthcare expenditures and indirect costs such as lost productivity and reduced quality of life.^{12–16} Therefore, a thorough investigation of risk factors for myopia is crucial for developing effective interventions in children, especially before the onset of myopia.

Several prospective cohort studies have evaluated potential independent risk factors for myopia onset among primary schoolchildren. These studies identified risk factors including age, sex, residential area, parental myopia, daily activities, refractive error, and ocular biometric parameters such as axial length (AL) and corneal refractive power.^{6–9,17–21} Routine clinical examinations typically include assessments of binocular vision to optimize visual clarity and minimize asthenopia, which should be managed if necessary. However, evidence regarding the role of binocular vision in myopia development and progression remains limited.²² In 1988, Goss et al. conducted a 3-year cohort study involving 87 emmetropic schoolchildren aged 8.5 to 14.3 years in the USA.^{23–24} Based on univariable analysis, they found that binocular visual function parameters such as positive relative accommodation (PRA), accommodative convergence to accommodation (AC/A) ratio, near lateral heterophoria, and near fusional vergence range might be informative for predicting myopia onset. However, it remains unclear whether these parameters remain significant after adjusting for confounders. Our previous report indicated that PRA could be a potential predictor of myopia onset over a relatively short follow-up period of 2 years,²⁵ but this observation requires verification over a longer follow-up period. Such information would be valuable for enhancing the accuracy of myopia prediction in the future.

Thus, the aim of this study was to identify independent risk factors for myopia onset, with a focus on exploring

various binocular visual function parameters, and to evaluate their potential value in predicting the occurrence of myopia.

Methods

Study design and population

The Wenzhou Medical University Essilor Progression and Onset of Myopia (WEPrOM) study was a school-based prospective cohort study conducted in Wenzhou, an eastern city in China. The study was approved by the ethics committee of the Eye Hospital of Wenzhou Medical University (KYK [2013]34) and adhered to the tenets of the Declaration of Helsinki.

The study involved two elementary schools in Wenzhou, one situated in the urban city center and the other in a rural area. During the baseline visit in November 2014, all 1118 second and third-grade students were invited to participate, with 1103 (98.7%) attending. We chose these two grades to focus on the critical ages for myopia onset while maximizing recruitment efficiency and follow-up rates. Of the 1103 screened children, 30 (2.7%) were excluded due to reported histories of ocular diseases affecting vision, such as strabismus, amblyopia, or congenital glaucoma. Consequently, 1073 schoolchildren were enrolled in the study. Follow-up visits were conducted annually until graduation at grade 6, except for the last two visits, which were delayed by six months due to a severe flu outbreak and the subsequent winter vacation. Written informed consent was obtained from the parents of all participants.

Measurements and outcomes

Trained investigators conducted comprehensive standardized ocular examinations at both schools using equipment calibrated at the start of each visit. Detailed descriptions of the measurement methods and outcomes are provided in the eMethods section of the Supplement. Specifically, we measured monocular uncorrected visual acuity (UCVA) and non-cycloplegic subjective refractive error for both eyes, as well as AL and corneal refractive power (CR) in the right

eye. Routine clinical binocular visual function parameters were also assessed, including near lateral heterophoria, AC/A ratio, negative relative accommodation (NRA), PRA, and the base-in (BI) and base-out (BO) break points of horizontal fusional convergence range at near. Prior to each ocular examination, questionnaires were distributed to parents to collect data on demographic characteristics (age, gender, school, and grade), daily activities (average time spent on near work and outdoor activities per day), parental myopia status (parental myopia, parental early-onset myopia [<12 years old], and parental high myopia), and parental educational attainment (both paternal and maternal). Refractive errors were calculated as spherical equivalent refraction (SER), defined as the sphere power plus half of the cylindrical power. Myopia was defined as an $SER \leq -0.50$ diopters (D) and further classified into low myopia ($-3.00 D < SER \leq -0.50 D$), moderate myopia ($-6.00 D < SER \leq -3.00 D$), and high myopia ($SER \leq -6.00 D$).^{7,26–27}

Statistical analysis

The right eye was selected for data analysis due to the high correlation in SER between the left and right eyes at each visit (Pearson correlation coefficient > 0.85 , $P < .001$ for all visits; see eTable 1 in the Supplement). Paired t -tests and McNemar tests were utilized for comparisons. To examine factors associated with myopia onset before primary school graduation, we conducted multivariable logistic regression analysis on participants who did not have myopia at baseline. Variables with P -values < 0.10 in the univariable analysis were included as potential confounders. We also adjusted for grade because follow-up durations differed between baseline grade 2 and grade 3 participants. Additionally, exploratory subgroup analyses were performed for significant parameters related to binocular visual functions.²⁸ To investigate possible modifications on the association between these parameters and myopia onset, we included demographic interaction terms in the logistic regression model and presented the results in a forest plot. P values for interaction were evaluated using interaction terms and likelihood ratio tests. To assess the predictive ability of factors associated with myopia onset before graduation, receiver operating characteristic (ROC) curves were plotted, and area under the curve (AUC) values were calculated. Statistical analyses were conducted using Stata/MP version 15.1 (Stata Corp LLC, TX, USA) and RStudio version 1.3 (The R Foundation for Statistical Computing, Vienna, Austria). Two-sided P values < 0.05 were considered statistically significant.

Results

After excluding 207 schoolchildren who were lost to follow-up due to reasons such as school transfer, absence from examinations, or the use of orthokeratology lenses or atropine, a total of 866 participants (mean [SD] baseline age: 7.8 [0.7] years; 400 [46.2%] female) with complete refraction data were included in the analysis. The characteristics of these 866 participants were compared between their baseline and graduation visits (Table 1). Compared to the baseline visit, significant increases were observed in age,

AL, AC/A ratio, and time spent on near work; moreover, SER values became significantly more negative. However, at the graduation visit, there were significant decreases in CR, the magnitude of NRA and PRA, BI break point, BO break point, and time spent outdoors.

Among the 714 participants without myopia at baseline, 444 (62.2%) developed myopia before graduation. After adjusting for confounders including baseline grade, children with the following baseline characteristics had a greater risk of myopia onset (Table 2): female sex (odds ratio [OR] = 3.03; $P < .001$), having two myopic parents (OR = 2.36; $P = .005$; compared to no myopic parents), worse UCVA (OR = 19.99; $P = .007$), more negative SER (OR = 0.15; $P < .001$), longer AL (OR = 7.28; $P < .001$), larger CR (OR = 2.20; $P < .001$), and lower magnitude of PRA (OR = 1.11; $P = .02$).

The correlation between baseline PRA and myopia onset before graduation remained consistent across all demographic subgroups (P -interaction > 0.05 for all; Figure 1). Consequently, the association between PRA and myopia onset did not exhibit significant variability. The ROC curves indicated that PRA had a moderate predictive capability for myopia onset before graduation, with an AUC of 0.59 (95% CI: 0.55 to 0.63), which was less effective compared to SER (AUC = 0.65; 95% CI: 0.61 to 0.69) but comparable to AL (AUC = 0.57; 95% CI: 0.53 to 0.62) and CR (AUC = 0.58; 95% CI: 0.53 to 0.62; Figure 2).

Discussion

Our data revealed that myopia onset before elementary school graduation was associated with sex, parental myopia, UCVA, SER values, AL, CR, and notably, the magnitude of PRA at baseline. The association between PRA and myopia onset remained robust across various demographic characteristics. Moreover, PRA demonstrated a predictive capability comparable to key ocular biometric parameters such as AL and CR.

Consistent with prior studies, we found that female sex,^{7,17} having two myopic parents,^{8–9,17–18,21} more myopic refractive error,^{6–7,9,19,21} longer AL,^{20–21} and larger corneal refractive power²¹ at baseline were independently associated with myopia onset before primary school graduation. In our study, the myopia incidence did not differ significantly between rural and urban schools after multivariable adjustment. This finding contrasts with the one-year result reported in Taipei but aligns with the 4-year follow-up findings in Beijing.^{9,29} Such consistency may be attributable to urbanization driven by China's rapid economic development. Contrary to previous reports,^{17,19} age was not independently associated with myopia onset in our study, potentially due to the narrow age range of participants. Outdoor activities and near work were significantly associated with myopia onset in some studies,^{9,18,21} but not consistently across all studies.^{6,8} Notably, the lack of a significant association between outdoor activities and near work with myopia onset has been predominantly observed in studies involving Chinese children, consistent with our findings. This phenomenon may be explained by the increased time spent on near work, reduced outdoor exposure, and limited variability in environmental factors resulting from intense academic competition among Chinese students.

Table 1 Characteristics of the Included Participants at Baseline and Graduation Visits^a.

| Characteristics ^b | Baseline (n = 866) | Graduation (n = 866) | P Value ^c |
|---------------------------------------------------------------|-------------------------------|-------------------------------|----------------------|
| Age, y | 7.8 (0.7) | 11.8 (0.5) | <0.001 |
| Gender, No. (%) | | | NA |
| Male | 466 (53.8) | 466 (53.8) | |
| Female | 400 (46.2) | 400 (46.2) | |
| School, No. (%) | | | NA |
| Rural | 219 (25.3) | 219 (25.3) | |
| Urban | 647 (74.7) | 647 (74.7) | |
| Baseline grade, No. (%) | | | NA |
| 2 | 436 (50.3) | 436 (50.3) | |
| 3 | 430 (49.7) | 430 (49.7) | |
| No. of myopic parents, No. (%) | | | NA |
| 0 | 315 (36.5) | 315 (36.5) | |
| 1 | 343 (39.8) | 343 (39.8) | |
| 2 | 204 (23.7) | 204 (23.7) | |
| Parental high myopia, No. (%) | | | NA |
| Neither | 736 (85.5) | 736 (85.5) | |
| Either | 125 (14.5) | 125 (14.5) | |
| Parental early-onset myopia, No. (%) | | | NA |
| Neither | 786 (91.5) | 786 (91.5) | |
| Either | 73 (8.5) | 73 (8.5) | |
| Paternal education, No. (%) | | | NA |
| High school or less | 464 (53.9) | 464 (53.9) | |
| College or beyond | 397 (46.1) | 397 (46.1) | |
| Maternal education, No. (%) | | | NA |
| High school or less | 456 (52.9) | 456 (52.9) | |
| College or beyond | 406 (47.1) | 406 (47.1) | |
| Uncorrected visual acuity, logMAR ^d | 0.00 (0.20) | — | NA |
| Spherical equivalent refraction, D | -0.14 (0.75) | -1.61 (1.75) | <0.001 |
| Axial length, mm | 23.10 (0.78) | 24.33 (1.01) | <0.001 |
| Corneal refractive power, D | 43.17 (1.42) | 43.11 (1.40) | <0.001 |
| Near lateral heterophoria, PD | -3.36 (5.18) | -3.58 (6.88) | .39 |
| AC/A ratio, PD/D | 1.77 (2.71) | 3.02 (2.63) | <0.001 |
| Negative relative accommodation, D | 2.73 (0.78) | 2.26 (0.62) | <0.001 |
| Positive relative accommodation, D | -3.66 (2.00) | -2.71 (1.38) | <0.001 |
| Base-in break point, PD | 22.55 (6.03) | 20.35 (5.86) | <0.001 |
| Base-out break point, PD | 25.75 (6.87) | 22.91 (7.30) | <0.001 |
| Time spent on near work, h/d ^e | 2.4 (2.3) ^f | 4.9 (2.7) ^f | <0.001 |
| Time spent outdoors, h/d ^e | 2.2 (1.9) ^f | 1.4 (1.0) ^f | <0.001 |
| Prevalence rate of myopia, No./total No. (%) [95% CI] | 152/866 (17.6) [15.0 to 20.1] | 596/866 (68.8) [65.7 to 71.9] | <0.001 |
| Prevalence rate of high myopia, No./total No. (%) [95% CI] | 0/866 (0.0) [0.0 to 0.0] | 16/866 (1.8) [1.0 to 2.7] | <0.001 |

Abbreviations: y, age in years; NA, not applicable; logMAR, logarithm of the minimum angle of resolution; D, diopter; PD, prism diopter; AC/A, accommodative convergence to accommodation; PD/D, prism diopter per diopter; h/d, hours per day; CI, confidence interval.

^a Data involved in the baseline visit were obtained in 2014 when children were in grade 2 or grade 3. Data from the graduation visit were collected in 2018 and 2019 successively when children were all in grade 6.

^b Data are presented as mean (standard deviation) unless otherwise indicated.

^c Statistical analyses were determined by paired *t*-test or McNemar test.

^d Uncorrected visual acuity was not evaluated among some myopic participants during follow-up visits; therefore, only baseline data are presented to avoid bias from non-random missing data.

^e Daily activities were assessed differently in the final visit in 2019 by obtaining details such as daily mobile phone usage time during weekdays.

^f All characteristics had a low rate (<7.0%) of missing data, except for time spent on near work (32.8% at baseline; 49.4% at graduation) and time spent outdoors (23.0% at baseline; 49.5% at graduation).

We identified that a lower magnitude of baseline PRA served as an independent risk factor for myopia onset. Our findings align with the 1996 study by Goss and colleagues, which reported that individuals who developed myopia

exhibited a lower magnitude of PRA.²³ This implies that a diminished capacity to sustain accommodative effort under prolonged binocular near-viewing conditions may be linked to myopia onset, potentially mediated through hyperopic

Table 2 Multivariable Logistic Regression Analysis of Factors Associated with Myopia Onset before Graduation^a.

| Baseline Characteristics ^b | No Myopia Onset (n = 270) | Myopia Onset (n = 444) | Unadjusted | | Adjusted | |
|---------------------------------------|---------------------------|------------------------|----------------------|---------|------------------------|---------|
| | | | OR (95% CI) | P Value | OR (95% CI) | P Value |
| Age, y | 7.8 (0.7) | 7.7 (0.7) | 0.80 (0.65 to 0.99) | .04 | 1.01 (0.70 to 1.47) | .95 |
| Gender, No. (%) | | | | | | |
| Male | 154 (57.0) | 217 (48.9) | 1 [Reference] | NA | 1 [Reference] | NA |
| Female | 116 (43.0) | 227 (51.1) | 1.39 (1.02 to 1.88) | .03 | 3.03 (1.99 to 4.62) | <.0001 |
| School, No. (%) | | | | | | |
| Rural | 80 (29.6) | 101 (22.7) | 1 [Reference] | NA | 1 [Reference] | NA |
| Urban | 190 (70.4) | 343 (77.3) | 1.43 (1.02 to 2.01) | .04 | 0.86 (0.51 to 1.45) | .57 |
| Grade, No. (%) | | | | | | |
| 2 | 127 (47.0) | 254 (57.2) | 1 [Reference] | NA | 1 [Reference] | NA |
| 3 | 143 (53.0) | 190 (42.8) | 0.66 (0.49 to 0.90) | .008 | 0.49 (0.28 to 0.86) | .01 |
| No. of myopic parents, No. (%) | | | | | | |
| 0 | 131 (48.9) | 139 (31.4) | 1 [Reference] | NA | 1 [Reference] | NA |
| 1 | 97 (36.2) | 184 (41.5) | 1.79 (1.27 to 2.52) | .001 | 1.44 (0.91 to 2.28) | .12 |
| 2 | 40 (14.9) | 120 (27.1) | 2.83 (1.84 to 4.35) | <.0001 | 2.36 (1.29 to 4.31) | .005 |
| Parental high myopia, No. (%) | | | | | | |
| Neither | 245 (91.4) | 372 (84.2) | 1 [Reference] | NA | 1 [Reference] | NA |
| Either | 23 (8.6) | 70 (15.8) | 2.00 (1.22 to 3.30) | .006 | 1.52 (0.82 to 2.82) | .19 |
| Parental early-onset myopia, No. (%) | | | | | | |
| Neither | 252 (94.4) | 405 (91.8) | 1 [Reference] | NA | --- | --- |
| Either | 15 (5.6) | 36 (8.2) | 1.49 (0.80 to 2.78) | .21 | --- | --- |
| Paternal education, No. (%) | | | | | | |
| High school or less | 148 (55.2) | 228 (51.6) | 1 [Reference] | NA | --- | --- |
| College or beyond | 120 (44.8) | 214 (48.4) | 1.16 (0.85 to 1.57) | .35 | --- | --- |
| Maternal education, No. (%) | | | | | | |
| High school or less | 153 (57.1) | 220 (49.7) | 1 [Reference] | NA | 1 [Reference] | NA |
| College or beyond | 115 (42.9) | 223 (50.3) | 1.35 (0.99 to 1.83) | .06 | 1.02 (0.66 to 1.59) | .91 |
| Uncorrected visual acuity, logMAR | -0.07 (0.10) | -0.06 (0.09) | 6.50 (1.21 to 34.84) | .03 | 19.99 (2.24 to 178.44) | .007 |
| Spherical equivalent refraction, D | 0.24 (0.53) | 0.02 (0.22) | 0.10 (0.05 to 0.19) | <.001 | 0.15 (0.07 to 0.31) | <.001 |
| Axial length, mm | 22.84 (0.71) | 23.03 (0.67) | 1.50 (1.20 to 1.88) | <.001 | 7.28 (4.30 to 12.31) | <.001 |
| Corneal refractive power, D | 42.90 (1.40) | 43.30 (1.40) | 1.19 (1.07 to 1.33) | .002 | 2.20 (1.75 to 2.76) | <.001 |
| Near lateral heterophoria, PD | -3.88 (4.97) | -3.51 (5.21) | 1.01 (0.99 to 1.05) | .35 | --- | --- |
| AC/A ratio, PD/D | 1.71 (2.88) | 1.82 (2.60) | 1.01 (0.96 to 1.07) | .63 | --- | --- |
| Negative relative accommodation, D | 2.68 (0.79) | 2.70 (0.74) | 1.03 (0.85 to 1.26) | .74 | --- | --- |
| Positive relative accommodation, D | -4.32 (2.33) | -3.62 (1.83) | 1.18 (1.10 to 1.28) | <.001 | 1.11 (1.02 to 1.22) | .02 |
| Base-in break point, PD | 22.81 (6.26) | 22.16 (5.73) | 0.98 (0.96 to 1.01) | .16 | --- | --- |
| Base-out break point, PD | 25.27 (6.86) | 25.33 (6.49) | 1.00 (0.98 to 1.03) | .90 | --- | --- |
| Time spent on near work, h/d | 2.4 (2.5) | 2.4 (2.3) | 1.01 (0.93 to 1.09) | .90 | --- | --- |
| Time spent outdoors, h/d | 2.1 (1.8) | 2.3 (2.0) | 1.06 (0.97 to 1.16) | .22 | --- | --- |

Abbreviations: OR, odds ratio; CI, confidence interval; y, age in years; NA, not applicable; logMAR, logarithm of the minimum angle of resolution; D, diopter; PD, prism diopter; AC/A, accommodative convergence to accommodation; PD/D, prism diopter per diopter; h/d, hours per day.

^a Multivariable logistic regression analysis was conducted among 714 participants without myopia at baseline, including variables with P values < 0.10 in the univariable analysis and adjusting for grade.

^b Data are presented as mean (standard deviation) unless otherwise indicated.

retinal defocus.^{30–31} Notably, subgroup analyses revealed a consistent association between reduced PRA and myopia onset across various demographic and familial factors, including age, gender, school, grade, parental myopia status, and parental educational attainment. These results underscore the stability of PRA as a risk factor for myopia onset and highlight its importance in myopia prediction models and future investigative efforts.

As one of the pioneering studies investigating the independent association between binocular visual functions and myopia, our study possesses several unique strengths, including a large sample size, a relatively long follow-up period, and a high follow-up rate (4.5-year follow-up rate of 80.7%). Nevertheless, this study also has certain limitations that warrant consideration. First, refraction measurements were conducted without cycloplegia, which might lead to an

overestimation of myopic refractive error. However, we employed subjective refraction, utilizing a working lens of +2.00 D following retinoscopy performed by an experienced ophthalmologist. Compared with non-cycloplegic autorefraction, this method minimizes the impact of accommodation while maintaining operational efficiency and enhancing examination compliance. Furthermore, it facilitates the assessment of binocular visual functions, demonstrating practical value in large-scale epidemiological studies. Second, some questionnaire data were incomplete (baseline missing data: 32.8% for near work; 23.0% for outdoor activities). Considering the potential recall bias inherent in questionnaire-based assessments and discrepancies between reports from parents and children,³² future studies should incorporate objective and portable wearable devices to improve data accuracy.³³




Figure 1 Forest Plot of Positive Relative Accommodation with Various Demographic Subgroups. The dashed vertical line represents the odds ratio for the overall study population. Abbreviations: OR, odds ratio; CI, confidence interval; PRA, positive relative accommodation; SD, standard deviation.




Figure 2 Receiver Operating Characteristic Curves of Risk Factors for Predicting Myopia Onset before Primary School Graduation. Abbreviations: ROC, receiver operating characteristic; AUC, area under the curve; UCVA, uncorrected visual acuity; SER, spherical equivalent refraction; PRA, positive relative accommodation; AL, axial length.

Conclusions

In conclusion, this school-based 4.5-year cohort study conducted in Wenzhou initially identifies PRA as a stable and independent risk factor for myopia onset prior to elementary school graduation, with predictive capability comparable to key ocular biometric parameters. This finding can inform future studies to enhance the accuracy of myopia prediction, thereby assisting eye care professionals and health administrators in making more informed decisions before implementing interventions in children.

Funding

This work was supported by the Scientific Research Project of Zhejiang Provincial Department of Education [grant number Y202457210], the International S&T Cooperation Program of China [grant number 2014DFA30940], and the Wenzhou Medical University-Essilor International Research Center [WEIRC, grant numbers 95013006, 95016010]. The funders had no role in the conceptualization, design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Consent for publication

Parental written informed consent were obtained from all enrolled participants.

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due to privacy or ethical restrictions but are available from the corresponding author on reasonable request.

Author contributions

All authors made substantial contributions to conception and design, acquisition of data, or analysis and interpretation of data; took part in drafting the article or revising it critically for important intellectual content; gave final approval of the version to be published; and agree to be accountable for all aspects of the work.

Declaration of competing interest

Drs. Yee Ling Wong, and Björn Drobe are employees of Essilor International, Singapore. No honorarium, grant, or other form of payment was given to anyone to produce the manuscript. The authors have no other potential conflicts of interest to disclose.

Acknowledgements

Not applicable.

Supplementary materials

Supplementary material associated with this article can be found in the online version at [doi:10.1016/j.optom.2025.100577](https://doi.org/10.1016/j.optom.2025.100577).

References

- Morgan I, Ohno-Matsui K, Myopia Saw S. *Lancet*. 2012;379(9827):1739–1748. [https://doi.org/10.1016/s0140-6736\(12\)60272-4](https://doi.org/10.1016/s0140-6736(12)60272-4).
- Rudnicka A, Kapetanakis V, Wathern A, et al. Global variations and time trends in the prevalence of childhood myopia, a systematic review and quantitative meta-analysis: implications for aetiology and early prevention. *Br J Ophthalmol*. 2016;100(7):882–890. <https://doi.org/10.1136/bjophthalmol-2015-307724>.
- Dong L, Kang Y, Li Y, Wei W, Jonas J. Prevalence and time trends of Myopia in children and adolescents in China: a systemic review and meta-analysis. *Retina*. 2020;40(3):399–411. <https://doi.org/10.1097/iae.0000000000002590>.
- Fan D, Lam D, Lam R, et al. Prevalence, incidence, and progression of myopia of school children in Hong Kong. *Invest Ophthalmol Vis Sci*. 2004;45(4):1071–1075. <https://doi.org/10.1167/iovs.03-1151>.

- Lam C, Edwards M, Millodot M, Goh W. A 2-year longitudinal study of myopia progression and optical component changes among Hong Kong schoolchildren. *Optom Vis Sci*. 1999;76(6):370–380. <https://doi.org/10.1097/00006324-199906000-00016>.
- Lin Z, Vasudevan B, Gao T, Zhou H, Ciuffreda K, Liang Y. Refractive change and incidence of myopia among rural Chinese children: the Handan Offspring Myopia Study. *Br J Ophthalmol*. 2021. <https://doi.org/10.1136/bjophthalmol-2020-317811>.
- Wang S, Guo Y, Liao C, et al. Incidence of and factors associated with myopia and high myopia in Chinese children, based on refraction without cycloplegia. *JAMA Ophthalmol*. 2018;136(9):1017–1024. <https://doi.org/10.1001/jamaophthalmol.2018.2658>.
- Ma Y, Zou H, Lin S, et al. Cohort study with 4-year follow-up of myopia and refractive parameters in primary schoolchildren in Baoshan District, Shanghai. *Clin Exp Ophthalmol*. 2018;46(8):861–872. <https://doi.org/10.1111/ceo.13195>.
- Tsai D, Fang S, Huang N, et al. Myopia development among young schoolchildren: the Myopia investigation Study in Taipei. *Invest Ophthalmol Vis Sci*. 2016;57(15):6852–6860. <https://doi.org/10.1167/iovs.16-20288>.
- Zhou W, Zhang Y, Li H, et al. Five-year progression of refractive errors and incidence of myopia in school-aged children in western China. *J Epidemiol*. 2016;26(7):386–395. <https://doi.org/10.2188/jea.JE20140258>.
- Zhao J, Mao J, Luo R, Li F, Munoz S, Ellwein L. The progression of refractive error in school-age children: shunyi district, China. *Am J Ophthalmol*. 2002;134(5):735–743. [https://doi.org/10.1016/s0002-9394\(02\)01689-6](https://doi.org/10.1016/s0002-9394(02)01689-6).
- Ikuno Y. Overview of the complications of high myopia. *Retina*. 2017;37(12):2347–2351. <https://doi.org/10.1097/iae.0000000000001489>.
- Lim M, Gazzard G, Sim E, Tong L, Saw S. Direct costs of myopia in Singapore. *Eye (Lond)*. 2009;23(5):1086–1089. <https://doi.org/10.1038/eye.2008.225>.
- Naidoo K, Fricke T, Frick K, et al. Potential lost productivity resulting from the global burden of myopia: systematic review, meta-analysis, and modeling. *Ophthalmology*. 2019;126(3):338–346. <https://doi.org/10.1016/j.ophtha.2018.10.029>.
- Rose K, Harper R, Tromans C, et al. Quality of life in myopia. *Br J Ophthalmol*. 2000;84(9):1031–1034. <https://doi.org/10.1136/bjo.84.9.1031>.
- Sankaridurg P, Tahhan N, Kandel H, et al. IMI impact of myopia. *Invest Ophthalmol Vis Sci*. 2021;62(5):2. <https://doi.org/10.1167/iovs.62.5.2>.
- Saw S, Shankar A, Tan S, et al. A cohort study of incident myopia in Singaporean children. *Invest Ophthalmol Vis Sci*. 2006;47(5):1839–1844. <https://doi.org/10.1167/iovs.05-1081>.
- French A, Morgan I, Mitchell P, Rose K. Risk factors for incident myopia in Australian schoolchildren: the Sydney adolescent vascular and eye study. *Ophthalmology*. 2013;120(10):2100–2108. <https://doi.org/10.1016/j.ophtha.2013.02.035>.
- Lin Z, Vasudevan B, Mao G, et al. The influence of near work on myopic refractive change in urban students in Beijing: a three-year follow-up report. *Graefes Arch Clin Exp Ophthalmol*. 2016;54(11):2247–2255. <https://doi.org/10.1007/s00417-016-3440-9>.
- Saw S, Tong L, Chua W, et al. Incidence and progression of myopia in Singaporean school children. *Invest Ophthalmol Vis Sci*. 2005;46(1):51–57. <https://doi.org/10.1167/iovs.04-0565>.
- Zadnik K, Sinnott L, Cotter S, et al. Prediction of juvenile-onset myopia. *JAMA Ophthalmol*. 2015;133(6):683–689. <https://doi.org/10.1001/jamaophthalmol.2015.0471>.
- Wolffsohn JS, Jong M, Smith 3rd EL, et al. IMI 2021 reports and digest - reflections on the implications for clinical practice. *Invest Ophthalmol Vis Sci*. Apr 28 2021;62(5):1. <https://doi.org/10.1167/iovs.62.5.1>.

23. Goss D, Jackson T. Clinical findings before the onset of myopia in youth: 2. Zone of clear single binocular vision. *Optom Vis Sci*. 1996;73(4):263–268. <https://doi.org/10.1097/00006324-199604000-00008>.
24. Goss D, Jackson T. Clinical findings before the onset of myopia in youth: 3. Heterophoria. *Optom Vis Sci*. 1996;73(4):269–278. <https://doi.org/10.1097/00006324-199604000-00009>.
25. Wong Y, Yuan Y, Su B, et al. Prediction of myopia onset with refractive error measured using non-cycloplegic subjective refraction: the WEPROM Study. *BMJ Open Ophthalmol*. 2021;6(1):e000628. <https://doi.org/10.1136/bmjophth-2020-000628>.
26. Mutti D, Hayes J, Mitchell G, et al. Refractive error, axial length, and relative peripheral refractive error before and after the onset of myopia. *Invest Ophthalmol Vis Sci*. 2007;48(6):2510–2519. <https://doi.org/10.1167/iops.06-0562>.
27. Flitcroft D, He M, Jonas J, et al. IML - defining and classifying myopia: a proposed set of standards for clinical and epidemiologic studies. *Invest Ophthalmol Vis Sci*. 2019;60(3):M20–M30. <https://doi.org/10.1167/iops.18-25957>.
28. Wright J, Williamson J, Whelton P, et al. A randomized trial of intensive versus standard blood-pressure control. *N Engl J Med*. 2015;373(22):2103–2116. <https://doi.org/10.1056/NEJMoa1511939>.
29. Guo Y, Liu L, Tang P, et al. Outdoor activity and myopia progression in 4-year follow-up of Chinese primary school children: the Beijing Children Eye Study. *PLoS One*. 2017;12(4):e0175921. <https://doi.org/10.1371/journal.pone.0175921>.
30. Gwiazda J, Thorn F, Held R. Accommodation, accommodative convergence, and response AC/A ratios before and at the onset of myopia in children. *Optom Vis Sci*. 2005;82(4):273–278. <https://doi.org/10.1097/01.opx.0000159363.07082.7d>.
31. Mutti D, Mitchell G, Jones-Jordan L, et al. The response AC/A ratio before and after the onset of myopia. *Invest Ophthalmol Vis Sci*. 2017;58(3):1594–1602. <https://doi.org/10.1167/iops.16-19093>.
32. Rah M, Mitchell G, Mutti D, Zadnik K. Levels of agreement between parents' and children's reports of near work. *Ophthalmic Epidemiol*. 2002;9(3):191–203. <https://doi.org/10.1076/opep.9.3.191.1514>.
33. Wen L, Cheng Q, Cao Y, et al. The Clouclip, a wearable device for measuring near-work and outdoor time: validation and comparison of objective measures with questionnaire estimates. *Acta Ophthalmol*. 2021. <https://doi.org/10.1111/aos.14785>.