

# How Could Contact Lens Wearers Be at Risk of *Acanthamoeba* Infection? A Review

Youhanna W. Ibrahim<sup>1,2</sup>, David L. Boase<sup>1</sup> and Ian A. Cree<sup>2,3</sup>

## ABSTRACT

Contact lens wear is highly influential on the incidence of ulcerative keratitis worldwide, particularly in developed countries. The association between *Acanthamoeba* keratitis and contact lens wear is firmly established; it may account for up to 95% of the reported cases. Before the popularisation of soft contact lens wear, *Acanthamoeba* keratitis was extremely rare. In 2000 it was estimated that the number of contact lens wearers worldwide was about 80 million, out of whom 33 million were in the United States and 90% of them wore hydrogel soft lenses. Contact lens-related problems depend on many factors, such as lens material, wearing modality, lens hygiene, type of lens-caring solution, the degree of compliance of the lens user with lens wear and care procedures, lens overwear, sleeping in lenses, rate of changing lenses, and lens case hygiene. This paper is a thorough review of the literature aiming to highlight the role of one of the main risk factors of infectious keratitis, contact lens wear, and also to show the responsibility of lens users in aggravating this risk. (J Optom 2009;2:60-66 ©2009 Spanish Council of Optometry)

**KEY WORDS:** *Acanthamoeba*; keratitis; contact lenses; contact lens wearer; lens overwear.

## RESUMEN

La utilización de lentes de contacto es un factor que influye sumamente en la incidencia de la queratitis ulcerosa, en particular en los países desarrollados. La asociación entre queratitis por *Acanthamoeba* y el uso de lentes de contacto está firmemente establecida; podría representar aproximadamente hasta un 95% de los casos registrados. Antes de la popularización del uso de lentes de contacto, la queratitis por *Acanthamoeba* era muy poco frecuente. En el año 2000 se estimó que, a nivel mundial, había unos 80 millones de usuarios de lentes de contacto, de los cuales 33 millones residían en los Estados Unidos y, de ellos, el 90% utilizaban lentes de contacto blandas de hidrogel. Los problemas derivados del uso de lentes de contacto dependen de otros muchos factores, como el material del que está hecha la lente, la modalidad de uso, las medidas de higiene empleadas, el tipo de solución utilizada para la limpieza y el mantenimiento de la lente, el grado de cumplimiento terapéutico del usuario de las lentes respecto a su utilización y a su limpieza/mantenimiento, el uso de las lentes durante periodos excesivamente prolongados, el dormir con las lentes de contacto puestas, la frecuencia con la que el usuario se cambia de lentes y la limpieza de la funda de las lentes de contacto. Este artículo presenta una revisión minuciosa de la literatura, con el objetivo de resaltar el papel de uno de los principales factores de riesgo de desarrollo de queratitis infecciosa, como es el uso de lentes de contacto,

From the <sup>1</sup>Department of Ophthalmology, Queen Alexandra Hospital, Portsmouth PO6 3LY. <sup>2</sup>Department of Pathology, Queen Alexandra Hospital, Portsmouth PO6 3LY. <sup>3</sup>Moorfields Eye Hospital, City Road, London EC1V 2PD. United Kingdom

Received: 4 September 2008

Revised: 1 March 2009

Accepted: 2 March 2009

Corresponding author: Dr. Youhanna William Ibrahim  
e-mail: youhannawilliam@hotmail.co.uk

y con el fin de poner de relieve que el propio usuario de las lentes es, en ocasiones, el responsable de agravar este riesgo.

(J Optom 2009;2:60-66 ©2009 Consejo General de Colegios de Ópticos-Optometristas de España)

**PALABRAS CLAVE:** *Acanthamoeba*; queratitis; lentes de contacto; usuario de lentes de contacto; uso excesivo de las lentes de contacto.

## INTRODUCTION

*Acanthamoeba* is a ubiquitous pathogen that can be found worldwide with different incidence rates, showing the ability to survive in very harsh environmental circumstances.<sup>1</sup> *Acanthamoeba* keratitis is a potentially blinding corneal infection<sup>2</sup> that may aggressively affect both eyes<sup>3,4</sup> with the possibility to recur after penetrating keratoplasty.<sup>4</sup> Contact lens wear remains the main risk factor in transmitting *Acanthamoeba* trophozoites and cysts to the cornea.<sup>5,6</sup> A 7-year 1997-2003 survey<sup>7</sup> showed an increase in the number of hospitalised patients due to contact lens-related corneal ulcers, which correlated with the increase in the number of lens wearers. *Acanthamoeba* has a great affinity for the attachment to the corneal epithelium<sup>8</sup> and to contact lenses.<sup>9</sup> The tendency for *Acanthamoeba* to adhere to surfaces is a key first step in the pathogenesis of *Acanthamoeba* keratitis, particularly in contact lens wearers. Contact lenses affect corneal epithelium integrity in two different ways: directly, through the associated fitting-related abrasions<sup>10</sup> and indirectly, by altering the normal physiological and metabolic cellular activities.<sup>11</sup> These later changes render the epithelial cells in a hypoxic status that eventually alters their integrity. Corneal oxygenation is significantly reduced during contact lens overwear, particularly for those who sleep in their lenses overnight.<sup>12</sup> However, patient's compliance and some basic hygienic standards can effectively minimise the risk of *Acanthamoeba* keratitis.<sup>13</sup>

## CONTACT LENS-RELATED PROBLEMS

### Contact Lens-Induced Trauma

**Direct Traumatic Effect of Lenses.** The corneal epithelium, with its tight junctions, creates an important barrier against *Acanthamoeba* invasion to the underlying corneal structures. Corneal epithelial cells are more resistant to the cytopathic effect of *Acanthamoeba* trophozoites than keratocytes.<sup>14</sup> Contact lens wear cause minor corneal abrasions, which is the key initial step for *Acanthamoeba* infection. Martinez et al.<sup>10</sup> suggested that corneal trauma was the crucial factor for *Acanthamoeba* infection, rather than immunosuppression. The adherence of the *Acanthamoeba* protozoon to an intact corneal epithelium without trauma did not lead, in animal models,

to the development of keratitis. Corneal abrasion was absolutely essential for the induction of *Acanthamoeba* keratitis in hamsters infected with contaminated contact lenses.<sup>15</sup>

Corneal epithelial defects make it possible for the *Acanthamoeba* protozoon to attach to the epithelium and to subsequently invade the rest of the underlying stromal layers. Corneal injury exposes protein sites known as mannose glycoproteins on the surface of injured corneas. The adhesion of the *Acanthamoeba* to the corneal epithelial cells is the result of mutual interaction between corresponding mannose-binding glycoproteins on the adjacent surface membranes.<sup>8,16,17</sup> Sugar inhibition assays<sup>16,18</sup> revealed how *Acanthamoeba* can selectively bind with high affinity to mannose saccharides and not to non-mannosylated neoglycoproteins, such as galactose, fucose, galactosamine or lactose. The corneal surface mannose receptors stimulate *Acanthamoeba* to secrete pathogenic proteases<sup>16</sup> which, in turn, induce epithelium apoptosis<sup>19</sup> and facilitate amoeba invasion to the underlying stroma. Alizadeh et al.<sup>20</sup> showed that contact lens wear exacerbated *Acanthamoeba* keratitis through the secretion of mannose-induced protease 133. In addition, the attachment of *Acanthamoeba* to corneal epithelial cells helps the phagocytosis and digestion of bacteria<sup>21</sup> that provide an important nutrition source for *Acanthamoeba*. However, Sharma et al.<sup>22</sup> found no difference between the adherence ability of *Acanthamoeba* to corneal epithelial cells of normal non-lens wearers and that of asymptomatic contact lens wearers.

**Indirect Traumatic Effect of Lenses.** Contact lens wear is usually associated with corneal epithelial hypoxia and hypercapnia, particularly when sleeping in lenses or when using lenses having low oxygen transmissibility.<sup>11</sup> Carbon dioxide accumulation alters the normal metabolic pathways, which leads to a series of micro-structural changes affecting all corneal layers, such as epithelial microcysts, depletion of epithelial glycogen storage, lactic acid accumulation, corneal acidosis, epithelial oedema,<sup>11</sup> decreased mitotic rate, increased central corneal thinning,<sup>23</sup> corneal hypoesthesia,<sup>24</sup> compromised junctional integrity, increased epithelial cells permeability,<sup>25</sup> increased cellular epithelial fragility, epithelial punctation, microscopic abrasion, sloughing of the epithelium and, eventually, corneal ulceration. In addition, changes in tear film thickness and stability<sup>26</sup> and alteration of the normal profile of conjunctival commensals<sup>27</sup> have been recognised. All these changes collectively breach the natural extra ocular protective mechanisms, rendering the cornea an easy target to a wide array of pathogens, including the *Acanthamoeba*.

#### Attachment of *Acanthamoeba* to Contact Lenses

Contact lenses serve as a vehicle for the harbouring, transmission and delivery of microorganisms to the eye. *Acanthamoeba* has a high affinity for contact lens surfaces, a property that plays an important role in the pathogenesis of *Acanthamoeba* keratitis and creates an actual threat to contact lens wearers.<sup>28</sup> The adherence of *Acanthamoeba* trophozoites and cysts to contact lenses can be detected and quantified using different methods and techniques.<sup>9,29-31</sup> The ability of *Acanthamoeba* to attach to contact lenses is influenced by several factors:

**Contact Lens Material, Ionicity, and Water Content.** The manufacturing material affects contact lens ability as a mechanical host allowing attachment and transfer of *Acanthamoeba* trophozoites or cysts onto the corneal surface. The incidence of *Acanthamoeba* is much lower with rigid lenses, as compared with the soft type. The lower incidence of *Acanthamoeba* keratitis found in the Netherlands<sup>32</sup> was attributed to the greater proportion of Dutch contact lens wearers that used rigid gas-permeable lenses. Rigid gas-permeable lenses were recommended to hospital staff members wearing lenses, with the aim of minimising the risk of infectious keratitis, due to the easy removal of the attached *Acanthamoeba* trophozoites and cysts from the surface of this type of contact lenses.<sup>33</sup> Kilvington et al.<sup>34</sup> showed that cyst attachment occurred only for soft lenses, but not for gas-permeable ones. However, a significant adherence of trophozoites was detected in the case of rigid gas-permeable lenses, as compared with soft contact ones<sup>35</sup>, with greater affinity for the silicone acrylate material of rigid lenses than for the fluoropolymer material of those same lenses.<sup>36</sup>

The greater affinity of *Acanthamoeba* trophozoites for silicone hydrogel lenses, as compared with conventional hydrogel lenses ( $P < 0.001$ ) was attributed to the attachment characteristics of the polymer of silicon type.<sup>37,38</sup> The attachment of *Acanthamoeba* trophozoites to different soft contact lens materials, such as polymacon, etafilcon A, lidofilcon A, and bufilcon A varied significantly, with the greatest adherence being observed for lidofilcon A and the least for the etafilcon A lenses.<sup>29</sup> Adherence of cysts and trophozoites was found to be higher for the non-ionic than for the ionic disposable lenses.<sup>28</sup> Simmons et al.<sup>39</sup> suggested that the attachment of *Acanthamoeba* was highly dependant on the ionic nature and the water content of soft contact lenses. Collectively, these reasons could explain why *Acanthamoeba* adherence is higher for disposable and extended-wear soft lenses than for the conventional soft daily and rigid lenses.

**Duration of Exposure and Protozoon Concentration.** Both cysts and trophozoites showed an immediate adherence to contact lenses, which was observed to happen within 10 seconds after exposure.<sup>36</sup> *Acanthamoeba* adherence to lens surface increased significantly for longer exposure durations and for higher concentrations of inoculum.<sup>29,40</sup> The higher water content of disposable soft contact lenses allows longer time of lens wear and, in turn, gives enough time of exposure for the lenses to be loaded with *Acanthamoeba*. This could add to the reasons why disposable soft contact lenses wearers are at greater risk of suffering from *Acanthamoeba* keratitis than those wearers of other types of contact lenses. However, Sharma et al.<sup>22</sup> found no difference in *Acanthamoeba* adherence to different contact lenses with increasing exposure time.

***Acanthamoeba*-Life Stage.** *Acanthamoeba* trophozoite shows a greater tendency to adhere to contact lenses, as compared with the cystic form.<sup>29,36,40-42</sup> In contrast to the cystic form, Sharma et al.<sup>36</sup> noticed more adherence of trophozoites to rigid gas-permeable lenses than to soft ones. Similarly, Kelly et al.<sup>29</sup> observed more preference of trophozoites to adhere

to rigid gas-permeable lenses and polymethylmethacrylate (PMMA) contact lenses compared with the cystic form, which showed non specific similar rates of adherence to a variety of lenses, such as rigid gas-permeable, PMMA, daily and disposable soft lenses.

**Lens Surface Deposits.** Attachment of trophozoites and cysts to contact lenses is highly influenced by the presence of protein deposits on the lens surface. Protein deposits on contact lens surface increase the adhesion of other bacterial microbes like *Pseudomonas aeruginosa*, on which *Acanthamoeba* feeds.<sup>43</sup> Protein and lipid deposition on lens surface is mediated by the chemical structure of the lens material and its water content. The high water content and the ionic material of some disposable soft lenses allow for more deposition of proteins, a fact that could explain the greater affinity of the *Acanthamoeba* protozoan for worn lenses than for unworn ones.<sup>39</sup> Jones et al.<sup>44</sup> reported significant deposition of low levels of lysozyme and high levels of lipid on silicone hydrogel contact lens materials, as compared with ionic contact lens materials. The adhesion of *Acanthamoeba* in unwashed worn versus unwashed unworn contact lenses showed a significantly lower adherence of *Acanthamoeba* to new lenses.<sup>40,41</sup> The serine protease subtilisin A enzyme used for protein removal from contact lenses has been found to have no cysticidal action even after 24 hours of exposure.<sup>45</sup> However, it could lower the number of protozoa attached to lens surface through protein removal.

**Mechanical Ways Used in Contact Lens Care.** While shaking showed no significant effect on adherence, a post-incubation wash using phosphate buffered saline decreased the number of adherent cysts and trophozoites.<sup>28</sup>

Several studies suggested that a good wash significantly decreased the adherence of trophozoites and cysts to the contact lens surface,<sup>40,42</sup> though one study<sup>46</sup> suggested that washing had no effect on either *Acanthamoeba* stage. Rinsing contact lenses in saline using the flow method was significantly more effective than the immersion technique in removing adherent *Acanthamoeba* trophozoites from rigid gas-permeable lenses.<sup>41</sup> Wiping, rinsing, and rubbing of contact lenses and lens cases with multipurpose disinfecting solutions dislodged adherent cysts and trophozoites and reduced the associated microbiological load.<sup>47</sup> Recent studies, showed that multipurpose contact lens solutions that employed a manual rub regime were more effective in removing adherent loosely-bound deposits<sup>48</sup> and different pathogenic microbes<sup>49</sup> from soft hydrogel lenses than rinsing or soaking alone.

**Associated Bacterial Organisms.** The contamination of lens care systems with bacteria is an essential association in the development of *Acanthamoeba* keratitis. The bacterial microorganisms that adhere to the surfaces of contact lenses provide a good medium that facilitates attachment, feeding, survival, and growth of *Acanthamoeba*. *Acanthamoeba* can easily attach and grow on a lens surface previously loaded with bacterial microorganisms. Gorlin et al.<sup>50</sup> found that about 50% of the eyes infected with *Acanthamoeba* had positive cultures for bacteria. Other study<sup>51</sup> showed that 85% of contact lens

systems infected with *Acanthamoeba* were contaminated with bacterial strains, mainly with the aerobic gram-negative bacilli *P. aeruginosa* and *Xanthomonas maltophilia*.

Other bacterial microorganisms, such as *Flavobacterium breve*, *P. paucimobilis*, *P. fluorescens*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Escherichia coli*, *Enterobacter agglomerans*, *Flavobacterium indolgenes*, *Salmonella enterica*, *Corynebacterium xerosis*, *Serratia marcescens* and *Klebsiella pneumoniae* were isolated in patients with *Acanthamoeba* keratitis.<sup>20,52-56</sup> Alizadeh et al.<sup>20</sup> showed that *Acanthamoeba* could secrete increased amounts of pathogenic mannose-induced protease 133 upon exposure to *C. xerosis*.

*Acanthamoeba* trophozoites and cysts could retain viable bacteria with human pathogenic potential.<sup>57</sup> Intra-*Acanthamoeba* detection, survival, growth, and multiplication of salmonellae<sup>56</sup> and *P. aeruginosa*<sup>58,59</sup> were reported, with the possibility of reisolating<sup>58</sup> *P. aeruginosa* from *Acanthamoeba* cysts. *P. aeruginosa* could significantly enhance *Acanthamoeba* trophozoite attachment to hydrogel contact lenses,<sup>60</sup> but not to silicone ones.<sup>37</sup> The combination of *P. aeruginosa* and *Acanthamoeba* was assumed to be selectively exclusive, causing potentially devastating ocular infections in contact lens wearers.<sup>33,61</sup> Sodium salicylate reduced trophozoite attachment to hydrogel lenses when inoculated with *P. aeruginosa*. This effect was attributed to the inhibition of bacterial biofilm-formation, interference with the biofilm-amoebal attachment, or modification of the lens surface.<sup>60</sup>

**Contact Lens Disinfecting Solutions.** The use of ineffective contact lens disinfecting solutions is strongly linked to the threat of *Acanthamoeba* infection in contact lens wearers. A 10-year survey (1994-2004)<sup>62</sup> showed that *Acanthamoeba* was isolated in contact lenses and contact lens disinfecting solutions in all cases of *Acanthamoeba* keratitis. The one-step 3% hydrogen peroxide and multipurpose solutions were found to be ineffective in killing *Acanthamoeba* cysts and trophozoites, as well as bacteria and fungi. However, in addition to the broad antimicrobial activity of multipurpose solutions, they were found to be capable of reducing the adherence capability of *Acanthamoeba* to contact lenses.<sup>63</sup> Opti-Free express multipurpose solution significantly reduced the adherence of trophozoites and cysts when used to clean, rinse, and soak soft contact lenses.<sup>64</sup> Complete Easy Rub multipurpose solution was effective in removing bacteria, fungi and *Acanthamoeba* from silicone hydrogel lenses.<sup>49</sup>

## PROBLEMS CAUSED BY CONTACT LENS WEARERS

### Contact Lens Overwear

Corneal overwear-related problems could develop in the long term for any type of contact lenses, including those designed for extended wear. The overwear problems were influenced by the rate of oxygen transmission and permeability through the lens material, lens thickness, lens type, wearing modalities, replacement schedule, repeated wear of disposable lenses, and overnight sleep in lenses. A lower incidence of microbial keratitis was reported for silicone hydrogel lenses with high oxygen permeability than for other soft lenses having low oxygen permeability used with an extended

wear scheme.<sup>65,66</sup> However, occasional pathophysiological problems, such as diffuse corneal infiltration,<sup>67</sup> development of mucin balls, superior epithelial arcuate lesions, contact lens papillary conjunctivitis, corneal erosions,<sup>11,68</sup> corneal dryness and discomfort,<sup>69</sup> central corneal thinning,<sup>70</sup> and thickened conjunctival epithelium due to increased metaplasia<sup>71</sup> were reported with the overwear of silicone hydrogel lenses with high oxygen permeability. A significantly higher risk of bacterial keratitis and a greater incidence of complications, such as limbal neovascularisation and corneal oedema, were reported in wearers whose daily wear time was higher than 12 hours.<sup>72</sup>

### Overnight Sleep in Different Types of Contact Lenses

The cornea gets its oxygen supply directly from the air when the eye is opened and from the surrounding blood vessels when it is closed. The new versions of rigid and soft contact lenses were designed to allow oxygen delivery to the cornea at an almost similar level under either opened or closed-eye conditions.<sup>73</sup> However, corneal hypoxia, subepithelial infiltrations, immune ring formation,<sup>12</sup> changes in corneal curvature, central corneal thinning,<sup>70</sup> alteration in the number of polymorphonuclear leukocytes, and variations in the level of different inflammatory mediators in the tear film<sup>74</sup> were reported upon wearing contact lenses for multiple sleep cycles. The results of various surveys<sup>75-78</sup> suggested that the overnight wear of contact lenses was the main cause of microbial keratitis, with a greater concern for the immunocompromised patients, where the risk of unusual infections was very high.<sup>79</sup>

The overnight wear-related corneal changes and the risk of ulcerative keratitis was found to be significantly dependant on the lens type. Overnight wear of rigid gas-permeable contact lenses was associated with higher levels of corneal hypoxia and epithelial oedema, as compared with soft lenses.<sup>25</sup> However, Graham et al.<sup>80</sup> stated that the severity of corneal swelling with rigid gas-permeable lenses was not a reliable predictor of ocular complications. No significant difference was recognised in the bacterial binding ability to the corneal epithelium between the overnight wear of either conventional or hyper-oxygen transmissible rigid contact lenses.<sup>81</sup> However, central corneal epithelial thinning with visual impairment<sup>82</sup> and the increased risk of bacterial<sup>83,84</sup> and *Acanthamoeba* keratitis<sup>84-89</sup> were reported with the overnight wear of orthokeratology contact lenses.

The overnight wear of extended wear lenses increased the risk of ulcerative keratitis,<sup>90</sup> with the possibility of conjunctival cytologic changes.<sup>71,75,76</sup> Schein et al.<sup>76</sup> mainly attributed the high risk of microbial keratitis among the users of extended wear lenses to the overnight wear, rather than to lens hygiene or lens type. However, Brennan<sup>91</sup> contradicted the common perception of optometrists in Western societies, who state that the use of extended wear lenses could increase the risk of microbial keratitis and the loss of vision. Brennan correlated the safety of the overnight wear to the high oxygen transmissibility of silicone hydrogel contact lenses. Other surveys<sup>92,93</sup> denied the occurrence of clinically serious events of microbial keratitis with daily disposable contact lenses.

Corneal hypoxic changes, such as epithelial oedema and microcysts were not recognised among the overnight wearers of extended wear lenses, with no significant difference in limbal redness between them and the non-lens users.<sup>94</sup> Kenyon et al.<sup>95</sup> suggested that neither the level of overnight corneal swelling nor the period between removals could influence the incidence or the severity of corneal problems of extended wear lenses.

### Non-Compliant Contact Lens Users

The compliance of contact lens wearers with the recommended lens care hygiene procedures is crucial to reduce the risk of serious infections. D'Aversa et al.<sup>96</sup> reviewed the medical records of 12 patients and found that substandard lens-care methods were used in 13 out of 14 (92.9%) eyes infected with *Acanthamoeba*. The use of tap water for the care of contact lenses was widely accepted as the main risk factor in *Acanthamoeba* infection.<sup>5,85,97-102</sup>

In the United Kingdom, 91% of the soft contact lens wearers and 94% of the rigid lens wearers avoided the disease by the complete avoidance of water and the use of powerful lens disinfecting solutions.<sup>5</sup> In the USA, the withdrawal of salt tablets from the market was responsible for the decrease in the incidence of *Acanthamoeba* keratitis in the mid 80s.<sup>98</sup> A recent study<sup>103</sup> showed that the Chicago-area tap water contained a highly virulent *Acanthamoeba* strain that was contributing to the increased incidence rate of *Acanthamoeba* keratitis in this area. Swimming, diving, showering or washing the face while wearing contact lenses was reported to cause *Acanthamoeba* keratitis.<sup>97,104-108</sup> The 50-fold increase in the risk of *Acanthamoeba* keratitis among disposable contact lens users was largely attributable to repeated wear of lenses, lack of disinfection, and use of saline and chlorine-based solutions.<sup>75</sup>

The compliance of contact lens users with the recommended care procedures is ineffective if these solutions do not manage to kill *Acanthamoeba*.<sup>6,105,109</sup> To avoid the persistent use of non-sterile solutions by non-compliant lens wearers, Moore<sup>13</sup> recommended heat disinfection of lenses—between 70 and 80°C for 10 minutes—and the use of 3% hydrogen peroxide for 2-3 hours, 0.001% thimerosal with edetate for 4 hours, 0.005% benzalkonium chloride with edetate for 4 hours, 0.001% chlorhexidine for 4 hours or 0.004% chlorhexidine for 1 hour. Better compliance of contact lens wearers was achieved with the introduction of multipurpose solutions. The multipurpose solutions replaced the need for an additional rinsing solution, offering a single solution for the cleaning, disinfection and contact lens storage.<sup>110</sup> The multipurpose solutions provided potent antimicrobial protection with less toxic and less allergenic effects.<sup>48,49,75</sup>

The non-compliance of contact lens users could occur in the case of deliberate reuse of daily disposable contact lenses, when wearing expired lenses without replacement or if using cheap contact lenses purchased from unlicensed vendors.<sup>111</sup> Old contact lenses could colonise more microorganisms due to the increased lens surface tear and wear-related scratches<sup>40</sup> or to the accumulated deposits.<sup>43-45</sup> The relationship between repeated use of daily disposable lenses

and risk of *Acanthamoeba*<sup>112,113</sup> and microbial keratitis<sup>93,114</sup> is well established.

Daily disposable lenses were designed for single use only, where a new sterile set should be opened every morning and discarded in the evening. This wear modality aimed to provide a great hygienic advantage, by avoiding the necessity and the cost of disinfecting solutions and storage cases. For hygienic purposes, daily disposable lenses were recommended for those lens wearers having jobs entailing a great potential risk of infection, such as hospital staff members.<sup>33</sup> Dart et al.<sup>115</sup> reported that vision loss was less likely to occur for daily disposable than for reusable soft lens wearers, though no significant reduction in the risk of microbial keratitis was found for users of daily disposable and silicone hydrogel lenses. However, the assumed lens wearers' compliance could sometimes divert the attention of professionals from considering *Acanthamoeba* infection in daily disposable contact lens wearers. A delay of 17 days before starting the anti-*Acanthamoeba* treatment was reported.<sup>116</sup>

Radford et al.<sup>75</sup> stated that the low care philosophy of daily disposable lens use has resulted in an absolute absence of care, and emphasised the importance of warning patients against the increased risk of infection upon reuse of daily disposable contact lenses. The non-compliance of contact lens users has been attributed to the convenience of using multipurpose or one-step solutions instead of using two-step hydrogen peroxide solutions. Financial savings achieved by reusing daily disposable contact lenses is another factor.<sup>117</sup>

## CONCLUSION

Contact lens wear is the main cause of ulcerative keratitis, which could get seriously complicated with corneal scarring and lead to permanent vision loss. The association between *Acanthamoeba* keratitis and contact lens wear is firmly established. Contact lenses have a great impact on corneal epithelium integrity. This, added to the greater affinity of *Acanthamoeba* to adhere to either corneal or lens surfaces, increase the risk of keratitis in contact lens wearers. Lens hygiene, lens care solutions, wearing modalities and the compliance of lens users are important factors in the lens-keratitis relationship. Every lens wearer should be aware of what the main risk factors are and, when given the routine instructions regarding lens fitting and care, they should also be provided with a thorough explanation of how contact lens misuse can seriously affect vision.

## REFERENCES

- Ibrahim YW, Boase DL, Cree IA. Factors affecting the epidemiology of *Acanthamoeba* keratitis. *Ophthalmic Epidemiol.* 2007;14:53-60.
- Awwad ST, Heilman M, Hogan RN, et al. Severe reactive ischemic posterior segment inflammation in *Acanthamoeba* keratitis: a new potentially blinding syndrome. *Ophthalmology.* 2007;114:313-320.
- Hassanlou M, Bhargava A, Hodge WG. Bilateral *Acanthamoeba* keratitis and treatment strategy based on lesion depth. *Can J Ophthalmol.* 2006;41:71-73.
- Rama P, Matuska S, Viganò M, Spinelli A, Paganoni G, Brancato R. Bilateral *Acanthamoeba* keratitis with late recurrence of the infection in a corneal graft: a case report. *Eur J Ophthalmol.* 2003;13:311-314.
- Radford CF, Lehmann OJ, Dart JK. *Acanthamoeba* keratitis: multicentre survey in England 1992-6. National *Acanthamoeba* Keratitis Study Group. *Br J Ophthalmol.* 1998;82:1387-1392.
- Butler TK, Males JJ, Robinson LP, et al. Six-year review of *Acanthamoeba* keratitis in New South Wales, Australia: 1997-2002. *Clin Experiment Ophthalmol.* 2005;33:41-46.
- Verhelst D, Koppen C, Van Looveren J, Meheus A, Tassignon MJ, the Belgian Keratitis Study Group. Contact lens-related corneal ulcers requiring hospitalization: a 7-year retrospective study in Belgium. *Acta Ophthalmol Scand.* 2006;84:522-526.
- Jaison PL, Cao Z, Panjwani N. Binding of *Acanthamoeba* to mannose-glycoproteins of corneal epithelium: effect of injury. *Curr Eye Res.* 1998;17:770-776.
- Beattie TK, Seal DV, Tomlinson A, McFadyen AK, Grimason AM. Determination of amoebicidal activities of multipurpose contact lens solutions by using a most probable number enumeration technique. *J Clin Microbiol.* 2003;41:2992-3000.
- Martinez AJ, Janitschke K. *Acanthamoeba*, an opportunistic microorganism: a review. *Infection.* 1985;13:251-256.
- Liesegang TJ. Physiologic changes of the cornea with contact lens wear. *CLAO J.* 2002;28:12-27.
- Klein P. Corneal immune ring as a complication of soft extended wear contact lens use. *Optom Vis Sci.* 1991;68:853-857.
- Moore MB. *Acanthamoeba* keratitis and contact lens wear: the patient is at fault. *Cornea.* 1990;9:S33-S35.
- Kinnear FB. *Acanthamoeba* pathogenicity for corneal cells. *J Infect.* 2004;49:310-316.
- Van Klink F, Alizadeh H, He YG, et al. Chinese hamster model of *Acanthamoeba* keratitis: role of contact lenses, trauma, and Langerhans cells. *Invest Ophthalmol Vis Sci.* 1993;34:1937-1944.
- Yang Z, Cao Z, Panjwani N. Pathogenesis of *Acanthamoeba* keratitis: carbohydrate-mediated host-parasite interactions. *Infect Immun.* 1997;65:439-445.
- Garate M, Alizadeh H, Neelam S, Niederkorn JY, Panjwani N. Oral immunization with *Acanthamoeba castellanii* mannose-binding protein ameliorates amoebic keratitis. *Infect Immun.* 2006;74:7032-7034.
- Cao Z, Jefferson DM, Panjwani N. Role of carbohydrate-mediated adherence in cytopathogenic mechanisms of *Acanthamoeba*. *J Biol Chem.* 1998;273:15838-15845.
- Zheng X, Uno T, Goto T, Zhang W, Hill JM, Ohashi Y. Pathogenic *Acanthamoeba* induces apoptosis of human corneal epithelial cells. *Jpn J Ophthalmol.* 2004;48:23-29.
- Alizadeh H, Neelam S, Hurt M, Niederkorn JY. Role of contact lens wear, bacterial flora, and mannose-induced pathogenic protease in the pathogenesis of amoebic keratitis. *Infect Immun.* 2005;73:1061-1068.
- Allen PG, Dawidowicz EA. Phagocytosis in *Acanthamoeba*. (I). A mannose receptor is responsible for the binding and phagocytosis of yeast. *J Cell Physiol.* 1990;145:508-513.
- Sharma S, Sankaridurg PR, Venkata N, et al. Adherence of *Acanthamoeba* to human corneal epithelial cells recovered from normal non-lens wearers and asymptomatic contact lens wearers. *Cont Lens Anterior Eye.* 1999;22:110-115.
- Myrowitz EH, Melia M, O'Brien TP. The relationship between long-term contact lens wear and corneal thickness. *CLAO J.* 2002;28:217-220.
- Ahuja M. Contact lens wear and microbial keratitis. *J Indian Med Assoc.* 2002;100:664-666.
- Lin MC, Graham AD, Fusaro RE, Polse KA. Impact of rigid gas-permeable contact lens extended wear on corneal epithelial barrier function. *Invest Ophthalmol Vis Sci.* 2002;43:1019-1024.
- Tutt T, Bradley A, Begley C, Thibos LN. Optical and visual impact of tear break-up in human eyes. *Invest Ophthalmol Vis Sci.* 2000;41:4117-4123.
- Barequet IS, Jabbur NS, Barron Y, Osterhout GJ, O'Brien TP. Perioperative microbiologic profile of the conjunctiva in photorefractive keratectomy. *J Refract Surg.* 2001;17:55-62.
- Ramachandran L, Janakiraman D, Sharma S, Rao GN. Effect of time and washing on the adhesion of *Acanthamoeba* to extended wear disposable hydrogel contact lenses. *CLAO J.* 1997;23:113-116.
- Kelly LD, Xu L. The effect of *Acanthamoeba* concentration on adherence to four types of unworn soft contact lenses. *CLAO J.* 1995;21:27-30.
- John T, Desai D, Sahm D. Adherence of *Acanthamoeba castellanii* cysts and trophozoites to unworn soft contact lenses. *Am J Ophthalmol.* 1989;108:658-664.
- Mathers WD, Sutphin JE, Folberg R, Meier PA, Wenzel RP, Elgin RG. Outbreak of keratitis presumed to be caused by *Acanthamoeba*. *Am J Ophthalmol.* 1996;121:129-142.
- Cheng KH, Leung SL, Hoekman HW, et al. Incidence of contact-lens-associated microbial keratitis and its related morbidity. *Lancet.* 1999;354:181-185.
- Hay J, Seal DV. Contact lens wear by hospital health care staff: is there cause for concern? *J Hosp Infect.* 1995;30:275-281.

34. Kilvington S, Larkin DF. *Acanthamoeba* adherence to contact lenses and removal by cleaning agents. *Eye*. 1990;4:589-590.
35. Cohen EJ, Fulton JC, Hoffman CJ, Rapuano CJ, Liabson PR. Trends in contact lens-associated corneal ulcers. *Cornea*. 1996;15:566-570.
36. Sharma S, Ramachandran L, Rao GN. Adherence of cysts and trophozoites of *Acanthamoeba* to unworn rigid gas permeable and soft contact lenses. *CLAO J*. 1995;21:247-451.
37. Beattie TK, Tomlinson A, McFadyen AK, Seal DV, Grimason AM. (II). Enhanced attachment of *Acanthamoeba* to extended-wear silicone hydrogel contact lenses: a new risk factor for infection? *Ophthalmology*. 2003;110:765-771.
38. Beattie TK, Tomlinson A, Seal DV. Surface treatment or material characteristic: the reason for the high level of *Acanthamoeba* attachment to silicone hydrogel contact lenses. *Eye Contact Lens*. 2003;29:S40-S43; discussion S57-S59, S192-S194.
39. Simmons PA, Tomlinson A, Connor R, Hay J, Seal DV. Effect of patient wear and extent of protein deposition on adsorption of *Acanthamoeba* to five types of hydrogel contact lenses. *Optom Vis Sci*. 1996;73:362-368.
40. Sehgal R, Saini J, Singh KD, Bharti HS. *Acanthamoeba* adherence to soft contact lens and human corneal stroma. *Indian J Pathol Microbiol*. 2002;45:63-67.
41. Cancrini G, Iori A, Mancino R. *Acanthamoeba* adherence to contact lenses, removal by rinsing procedures, and survival to some ophthalmic products. *Parassitologia*. 1998;40:275-278.
42. Seal DV, Hay J, Kirkness CM. Chlorhexidine or polyhexamethylene biguanide for *Acanthamoeba* keratitis. *Lancet*. 1995;345:136.
43. Butrus SL, Klotz SA. Contact lens surface deposits increase the adhesion of *Pseudomonas aeruginosa*. *Curr Eye Res*. 1990;9:717-724.
44. Jones L, Senchyna M, Glasier MA, et al. Lysozyme and lipid deposition on silicone hydrogel contact lens materials. *Eye Contact Lens*. 2003;29:S75-S79; discussion S83-S84, S192-S194.
45. Hughes R, Kilvington S. Comparison of hydrogen peroxide contact lens disinfection systems and solutions against *Acanthamoeba* polyphaga. *Antimicrob Ag Chemother*. 2001;45:2038-2043.
46. John T. Interactions of bacteria and amoebae with ocular biomaterials. *Cells and Materials*. 1991;1:129-139.
47. Niszl IA, Markus MB. Treatment of *Acanthamoeba* keratitis. *S Afr Med J*. 1996;86:566.
48. Cho P, Cheng SY, Chan WY, Yip WK. Soft contact lens cleaning: rub or no-rub? *Ophthalmic Physiol Opt*. 2009;29:49-57.
49. Kilvington S, Lonnen J. A comparison of regimen methods for the removal and inactivation of bacteria, fungi and *Acanthamoeba* from two types of silicone hydrogel lenses. *Cont Lens Anterior Eye*. 2009;32:73-77.
50. Gorlin AI, Gabriel MM, Wilson LA, Ahearn DG. Effect of adhered bacteria on the binding of *Acanthamoeba* to hydrogel lenses. *Arch Ophthalmol*. 1996;114:576-580.
51. Kelly LD, Xu L. The effect of concurrent *Pseudomonas* or *Xanthomonas* exposure on adherence of *Acanthamoeba castellanii* to soft contact lenses. *Graefes Arch Clin Exp Ophthalmol*. 1996;234:311-314.
52. Bottone EJ, Madayag RM, Qureshi MN. *Acanthamoeba* keratitis. Synergy between amoebic and bacterial co-contaminants in contact lens care systems as a prelude to infection. *J Clin Microbiol*. 1992;30:2447-2450.
53. Clark BJ, Harkins LS, Munro FA, Devonshire P. Microbial contamination of cases used for storing contact lenses. *J Infect*. 1994;28:293-304.
54. Wang X, Ahearn DG. Effect of bacteria survival and growth of *Acanthamoeba castellanii*. *Curr Microbiol*. 1997;34:212-215.
55. Willcox MD, Low R, Hon J, Harmis N. Does *Acanthamoeba* protect *Pseudomonas aeruginosa* from the bactericidal effects of contact lens disinfecting systems? *Aust N Z J Ophthalmol*. 1998;26:S32-S35.
56. Tezcan-Merdol D, Ljungstrom M, Winiacka-Krusnell J, Linder E, Engstrand L, Rhen M. Uptake and replication of *Salmonella enterica* in *Acanthamoeba* rhyodes. *Appl Environ Microbiol*. 2004;70:3706-3714.
57. Marciano-Cabral F, Cabral G. *Acanthamoeba* spp. as agents of disease in humans. *Clin Microbiol Rev*. 2003;16:273-307.
58. Walochnik J, Picher O, Aspöck C, Ullmann M, Sommer R, Aspöck H. Interactions of "Limax amoebae" and gram-negative bacteria: experimental studies and review of current problems. *Tokai J Exp Clin Med*. 1999;23:273-278.
59. Michel R, Burghardt H, Bergmann H. *Acanthamoeba*, naturally intracellularly infected with *Pseudomonas aeruginosa*, after their isolation from a microbiologically contaminated drinking water system in a hospital. *Zentralbl Hyg Umweltmed*. 1995;196:532-544.
60. Tomlinson A, Simmons PA, Seal DV, McFadyen AK. Salicylate inhibition of *Acanthamoeba* attachment to contact lenses: a model to reduce risk of infection. *Ophthalmology*. 2000;107:112-117.
61. Dini LA, Cockinos C, Frean JA, Niszl IA, Markus MB. Unusual case of *Acanthamoeba* polyphaga and *Pseudomonas aeruginosa* keratitis in a contact lens wearer from Gauteng, South Africa. *J Clin Microbiol*. 2000;38:826-829.
62. Tzanetou K, Miltakakis D, Droutsas D, et al. *Acanthamoeba* keratitis and contact lens disinfecting solutions. *Ophthalmologica*. 2006;220:238-241.
63. Borazjani RN, Kilvington S. Effect of a multipurpose contact lens solution on the survival and binding of *Acanthamoeba* species on contact lenses examined with a no-rub regimen. *Eye Contact Lens*. 2005;31:39-45.
64. Kilvington S, Anger C. A comparison of cyst age and assay method of the efficacy of contact lens disinfectants against *Acanthamoeba*. *Br J Ophthalmol*. 2001;85:336-340.
65. Holden BA, Sweeney DF, Sankaridurg PR, et al. Microbial keratitis and vision loss with contact lenses. *Eye Contact Lens*. 2003;29:S131-S134; discussion S143-S144, S192-S194.
66. Bialasiewicz AA. Infection immunology in silicone hydrogel contact lenses for continuous wear--a review. *Klin Monatsbl Augenheilkd*. 2003;220:453-458.
67. Skotnitsky C, Jalbert I, O'Hare N, Sweeney DF, Holden BA. Case reports of three atypical infiltrative keratitis events with high Dk soft contact lens wear. *Cornea*. 2002;21:318-324.
68. Dumbleton K. Noninflammatory silicone hydrogel contact lens complications. *Eye Contact Lens*. 2003;29:S186-S189; discussion S190-S191, S192-S194.
69. Fonn D, Dumbleton K. Dryness and discomfort with silicone hydrogel contact lenses. *Eye Contact Lens*. 2003;29:S101-S104; discussion S115-S118, S192-S194.
70. Gonzalez-Meijome JM, Gonzalez-Perez J, Cervino A, Yebra-Pimentel E, Parafita MA. Changes in corneal structure with continuous wear of high-Dk soft contact lenses: a pilot study. *Optom Vis Sci*. 2003;80:440-446.
71. Gurdal C, Aydin S, Kirimlioglu H, Toprak E, Sengor T. Effects of extended-wear soft contact lenses on the ocular surface and central corneal thickness. *Ophthalmologica*. 2003;217:329-336.
72. Rapkin JS. The effect of daily wear time on contact lens complications. *CLAO J*. 1988;14:139-142.
73. Weissman B, Mondino BJ. Why daily wear is still better than extended wear. *Eye Contact Lens*. 2003;29:S145-S146.
74. Thakur A, Willcox MD. Contact lens wear alters the production of certain inflammatory mediators in tears. *Exp Eye Res*. 2000;70:255-259.
75. Radford CF, Bacon AS, Dart JK, Minassian DC. Risk factors for *Acanthamoeba* keratitis in contact lens users: a case-control study. *BMJ*. 1995;310:1567-1570.
76. Schein OD, Buehler PO, Stamler JF, Verdier DD, Katz J. The impact of overnight wear on the risk of contact lens-associated ulcerative keratitis. *Arch Ophthalmol*. 1994;112:186-190.
77. Lam DS, Houang E, Fan DS, Lyon D, Seal D, Wong E. Incidence and risk factors for microbial keratitis in Hong Kong: comparison with Europe and North America. *Eye*. 2002;16:608-618.
78. Landers JA, Crompton JL. Microbial keratitis associated with overnight wear of silicone hydrogel contact lenses. *Med J Aust*. 2006;185:177-178.
79. Sridhar MS, Laibson PR, Rapuano CJ, Cohen EJ. Infectious crystalline keratopathy in an immunosuppressed patient. *CLAO J*. 2001;27:108-110.
80. Graham AD, Fusaro RE, Polse KA, Lin MC, Giasson CJ. Predicting extended wear complications from overnight corneal swelling. *Invest Ophthalmol Vis Sci*. 2001;42:3150-3157.
81. Cavanagh HD, Ladage P, Yamamoto K, Li SL, Petroll WM, Jester JV. Effects of daily and overnight wear of hyper-oxygen transmissible rigid and silicone hydrogel lenses on bacterial binding to the corneal epithelium: 13-month clinical trials. *Eye Contact Lens*. 2003;29:S14-S16; discussion S26-S29, S192-S194.
82. Alharbi A, Swarbrick HA. The effects of overnight orthokeratology lens wear on corneal thickness. *Invest Ophthalmol Vis Sci*. 2003;44:2518-2523.
83. Lau LI, Wu CC, Lee SM, Hsu WM. *Pseudomonas* corneal ulcer related to overnight orthokeratology. *Cornea*. 2003;22:262-264.
84. Sun X, Zhao H, Deng S, et al. Infectious keratitis related to orthokeratology. *Ophthalmic Physiol Opt*. 2006;26:133-136.
85. Watt K, Swarbrick HA. Microbial keratitis in overnight orthokeratology: review of the first 50 cases. *Eye Contact Lens*. 2005;31:201-208.
86. Hutchinson K, Apel A. Infectious keratitis in orthokeratology. *Clin Experiment Ophthalmol*. 2002;30:49-51.

87. Lu L, Zou L, Wang R. Orthokeratology induced infective corneal ulcer. *Zhonghua Yan Ke Za Zhi*. 2001;37:443-446.
88. Xuguang S, Lin C, Yan Z, et al. Acanthamoeba keratitis as a complication of orthokeratology. *Am J Ophthalmol*. 2003;136:1159-1161.
89. Wilhelmus KR. Acanthamoeba keratitis during orthokeratology. *Cornea*. 2005;24:864-866.
90. Schein OD, Poggio EC. Ulcerative keratitis in contact lens wearers. Incidence and risk factors. *Cornea*. 1990;9:S55-S58; discussion S62-S63.
91. Brennan NA. Is there a question of safety with continuous wear? *Clin Exp Optom*. 2002;85:127-140.
92. Levy B, McNamara N, Corzine J, Abbott RL. Prospective trial of daily and extended wear disposable contact lenses. *Cornea*. 1997;16:274-276. Erratum in: *Cornea* 1997;16:600.
93. Sankaridurg PR, Sweeney DF, Holden BA, et al. Comparison of adverse events with daily disposable hydrogels and spectacle wear: results from a 12-month prospective clinical trial. *Ophthalmology*. 2003;110:2327-2334.
94. Sweeney DF. Clinical signs of hypoxia with high-Dk soft lens extended wear: is the cornea convinced? *Eye Contact Lens*. 2003;29:S22-S25; discussion S26-S29, S192-S194.
95. Kenyon E, Polse KA, Seger RG. Influence of wearing schedule on extended-wear complications. *Ophthalmology*. 1986;93:231-236.
96. D'Aversa G, Stern GA, Driebe WT Jr. Diagnosis and successful medical treatment of Acanthamoeba keratitis. *Arch Ophthalmol*. 1995;113:1120-1123.
97. Seal DV, Dalton A, Doris D. Disinfection of contact lenses without tap water rinsing-is it effective? *Eye*. 1999;13:226-230.
98. Mathers WD, Sutphin JE, Lane JA, Folberg R. Correlation between surface water contamination with amoeba and the onset of symptoms and diagnosis of amoeba-like keratitis. *Br J Ophthalmol*. 1998;82:1143-1146.
99. Jeong HJ, Yu HS. The role of domestic tap water in Acanthamoeba contamination in contact lens storage cases in Korea. *Korean J Parasitol*. 2005;43:47-50.
100. Visvesvara G.S. Epidemiology of infections with free-living amoebae and laboratory diagnosis of microsporidiosis. *Mt. Sinai J. Med*. 1993;60:283-288.
101. Cohen EJ, Buchanan HW, Laughrea PA, et al. Diagnosis and management of Acanthamoeba keratitis. *Am. J. Ophthalmol*. 1985;100:389-395.
102. Moore MB, McCulley JP, Newton C, Cobo LM, Foulks GN, O'Day DM. Acanthamoeba keratitis. A growing problem in soft and hard contact lens wearers. *Ophthalmology*. 1987;94:1654-1661.
103. Shoff ME, Joslin CE, Tu EY, Kubatko L, Fuerst PA. Efficacy of contact lens systems against recent clinical and tap water Acanthamoeba isolates. *Cornea*. 2008;27:713-719.
104. Radford CF, Minassian DC, Dart JK. Acanthamoeba keratitis in England and Wales: incidence, outcome, and risk factors. *Br J Ophthalmol*. 2002;86:536-542.
105. Stehr-Green JK, Bailey TM, Brandt FH, Carr CH, Bond WW, Visvesvara GS. Acanthamoeba keratitis in soft contact lens wearers: a case-control study. *JAMA*. 1987;258:57-60.
106. Ondriska F, Mrva M, Lichvar M, Ziak P, Murgasova Z, Nohynkova E. First cases of Acanthamoeba keratitis in Slovakia. *Ann Agric Environ Med*. 2004;11:335-341.
107. De Jonckheere JF. Epidemiological typing of Acanthamoeba strains isolated from keratitis cases in Belgium. *Bull Soc Belge Ophthalmol*. 2003;287:27-33.
108. Kaji Y, Hu B, Kawana K, Oshika T. Swimming with soft contact lenses: danger of Acanthamoeba keratitis. *Lancet Infect Dis*. 2005;5:392.
109. Niszl IA, Markus MB. Anti-Acanthamoeba activity of contact lens solutions. *Br J Ophthalmol*. 1998;82:1033-1038.
110. Stevenson RW, Seal DV. Has the introduction of multi-purpose solutions contributed to a reduction in Acanthamoeba keratitis in contact lens wearers? *Contact Lens Anterior Eye*. 1998;21:89-92.
111. Gagnon MR, Walter KA. A case of Acanthamoeba keratitis as a result of a cosmetic contact lens. *Eye Contact Lens*. 2006;32:37-38.
112. Blades K, Tomlinson A, Seal DV. Acanthamoeba keratitis occurring with daily disposable contact lens wear. *Br J Ophthalmol*. 2000;84:805.
113. Woodruff SA, Dart JK. Acanthamoeba keratitis occurring with daily disposable contact lens wear. *Br J Ophthalmol*. 1999;83:1088-1089.
114. Stapleton F. Contact lens-related microbial keratitis: what can epidemiologic studies tell us? *Eye Contact Lens*. 2003;29:S85-S89; discussion S115-S118, S192-S194.
115. Dart JK, Radford CF, Minassian D, Verma S, Stapleton F. Risk factors for microbial keratitis with contemporary contact lenses: a case-control study. *Ophthalmology*. 2008;115:1647-1654.
116. Niyadurupola N, Illingworth CD. Acanthamoeba keratitis associated with misuse of daily disposable contact lenses. *Cont Lens Anterior Eye*. 2006;29:269-271.
117. Hiti K, Walochnik J, Haller-Schober EM, Faschinger C, Aspöck H. Viability of Acanthamoeba after exposure to a multipurpose disinfecting contact lens solution and two hydrogen peroxide systems. *Br J Ophthalmol*. 2002;86:144-146.