



## ORIGINAL ARTICLE

## Convergence insufficiency symptom survey (CISS) scores are predictive of severity and number of clinical signs of convergence insufficiency in young adult Africans

Charles Darko-Takyi<sup>a,\*</sup>, Andrew Owusu-Ansah<sup>a</sup>, Frank Boampong<sup>b</sup>, Enyam Komla Morny<sup>a</sup>, Francisca Hammond<sup>a</sup>, Stephen Ocansey<sup>a</sup>

<sup>a</sup> Department of Optometry and Vision Science, University of Cape Coast, Ghana

<sup>b</sup> Methodist Hospital Eye Clinic, Kumasi, Ghana

Received 27 November 2020; accepted 4 May 2021

Available online 18 October 2021



### KEYWORDS

Correlations;  
Sensitivity;  
Specificity;  
Convergence insufficiency symptoms survey score;  
Convergence insufficiency;  
University students

### Abstract

**Purpose:** This study aimed to investigate the correlation between convergence insufficiency symptom survey (CISS) score and the signs of convergence insufficiency (CI) and to evaluate the sensitivity and specificity of the CISS to predict CI and ametropia among young adult university students.

**Method:** This prospective cross-sectional clinic-based study included 300 first year university students (mean age = 21.58 (SD ± 2.2) years) who consecutively reported for eye examination. Participants were administered the CISS questionnaire and investigated for the signs of CI. Diagnosis of CI was based on presence of three or four signs. The correlation between the CISS score and the signs of CI were determined and Receiver Operation Characteristics (ROC) curves were used to evaluate sensitivity and specificity.

**Results:** There were significant correlations between CISS score and the clinical signs of CI namely NPC break ( $r_s = 0.622$ ,  $p = 0.0001$ ), NPC recovery ( $r_s = 0.620$ ,  $p = 0.0001$ ), near exophoria ( $r_s = 0.434$ ,  $p = 0.0001$ ), near PFV blur ( $r_s = -0.359$ ,  $p = 0.0001$ ), near PFV break ( $r_s = -0.306$ ,  $p = 0.0001$ ), near PFV recovery ( $r_s = -0.326$ ,  $p = 0.0001$ ) and gradient AC/A ratio ( $r_s = -0.290$ ,  $p = 0.0001$ ). There was a significant positive correlation between CISS score and the number of clinical signs of CI ( $r_s = 0.575$ ,  $p$ -value = 0.0001). The CISS had good sensitivity (AOC = 0.882) to predict CI and poor sensitivity (AOC = 0.642) to predict ametropia.

**Conclusion:** The CISS score is correlated with the severity and number of signs of CI in young adult Ghanaian university students. Its use in addition to clinical investigative testing may give a definitive diagnosis of symptomatic CI.

© 2021 Spanish General Council of Optometry. Published by Elsevier España, S.L.U. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

\* Corresponding author.

E-mail address: [cdarko-takyi@ucc.edu.gh](mailto:cdarko-takyi@ucc.edu.gh) (C. Darko-Takyi).

### Abbreviations

CI	Convergence insufficiency
CISS	Convergence insufficiency symptom survey
ROC	Receiver Operating Characteristics
AOC	Area under the curve
PFV	Positive fusional vergence
NPC	Near point of convergence

## Introduction

Convergence insufficiency (CI) is the commonest vergence disorder in young adult African populations.<sup>1,2</sup> It is typically characterized by symptoms of asthenopia mostly during reading or performing near work and clinical signs such as exophoria greater at near than distance, receded near point of convergence, decreased positive fusional vergence at near and other signs of low accommodative convergence over accommodation (AC/A) ratio.<sup>3</sup> Recent reported prevalence's among African university students range from 1.6% among Optometry students in Ghana<sup>2</sup> to 29.6% in first year University students in Benin City, Nigeria.<sup>1</sup> A specialized questionnaire for CI (Convergence Insufficiency Symptom Survey, CISS) has been shown to have good construct validity and reliability and has been used to measure the type, severity and frequency of symptoms before and after treatment for patients with symptomatic CI.<sup>4,5</sup> There is evidence of a case of CI in an African patient who was successfully treated using the CISS scores as guide to evaluate outcome.<sup>6</sup> In a recent study, it has been confirmed that examiner bias in terms of CISS questionnaire administration does not affect CISS scores.<sup>7</sup>

Rouse et al.<sup>8</sup> in their study among 8 to 12 year old children suggested a potential correlation between patient symptoms and the number of clinical signs of CI even though there was no statistically reported findings to support this assertion. Another study among symptomatic children with a CISS score greater than 16 and three clinical signs of CI, found no correlation between the severity of the clinical signs and their level of symptoms.<sup>9</sup> One study, however, determined significant weak correlations between symptoms and clinical signs of CI among children 8 to 10 years of age.<sup>10</sup> This latter study<sup>10</sup>, however, did not administer the CISS. Study by Horwood et al.<sup>11</sup> on young adults showed that a high CISS score can be a good predictor of presence of one or more signs of CI. However, authors did not find any study that has investigated specific correlations between the clinical signs of CI and level of symptoms among young adults. With different CISS cut off points for symptomatic and asymptomatic young adults<sup>12</sup> and children,<sup>13</sup> it is likely that the findings on relationship between signs and symptom levels might differ.

A study indicated that the CISS is not specific for CI;<sup>14</sup> the originators of the revised CISS also acknowledge that it could be helpful in patients with other accommodative or vergence disorders<sup>13</sup> because their symptoms are similar.<sup>3</sup> Borsting et al.<sup>15</sup> also found out that the CISS was able to discriminate between children with accommodative insufficiency and normal binocular vision. With ametropia been one of the major conditions encountered in the optometric

clinic population,<sup>16</sup> it was prudent to investigate possible associations between CISS score and ametropia. The symptoms of some types of ametropia and accommodative or binocular vision anomalies may overlap.<sup>17</sup>

The present study aimed to investigate the correlation between CISS score, and severity and number of clinical signs of CI among a young adult African University population. It also aimed to evaluate the sensitivity of the CISS to predict CI among these young adult African students. The present study also sought to explore possible associations between CISS scores and magnitudes of ametropia and to evaluate the sensitivity and specificity of the CISS to predict ametropia among these young adult African students.

## Methodology

### Ethical considerations

This clinic-based prospective cross sectional study adhered to the tenets of the declaration of Helsinki. Institutional ethical approval was obtained from Institutional Review Board of University of Cape Coast (Reference number: UCCIRB/CHAS/2015/057). The participants signed a written informed consent before enrolling each of them into the study.

### Study participants, setting and sampling procedure

The study was conducted in the optometric clinic situated on the campus of the University of Cape Coast, Ghana. The study population were first year university students (admitted within the 2015/2016 academic year) who reported to the optometric clinic for the university's mandatory first year student's eye examination. All first year students who consecutively attended the clinic for eye examination within the study period of October, 2015 to April, 2016 were enrolled into the study if they gave informed consent.

## Clinical procedures

### Visual acuity

Distance visual acuity (VA) at 6 m and near VA at 0.4 m was taken using a distance Snellen chart and a near N-notation visual acuity chart respectively.

### External and internal eye examination

The ocular adnexa and anterior segment of both eyes were examined under a slit lamp biomicroscope. (Topcon SL-2G); the posterior segments of both eyes were examined using Keeler professional direct ophthalmoscope.

### Inclusion / eligibility criteria

First year students aged 18 to 30 years old were included in the study. Students with ocular diseases, strabismus, nystagmus, unilateral or bilateral blindness, or on medications or any form of medical treatment were excluded from the study.

## Questionnaire administration

The 15 item revised CISS questionnaire (with each item scored on a scale of 0 to 4, where 4 represents the highest frequency of occurrence of symptoms "always" and 0 represents no symptom)<sup>12</sup> with a good internal consistency (Chronbach alpha = 0.880) among Ghanaian University students, was administered to selected participants. All 15 item scores for each participant were summed to obtain the total CISS score. A CISS score greater than or equal to 21 was considered symptomatic.<sup>12</sup>

## Refraction

Objective refraction was determined with static retinoscopy (Keeler Professional streak retinoscope) followed by subjective refraction with a manual phoropter (Topcon VT-10).<sup>18</sup> Final refractive correction was determined as the highest plus lens or least minus lens for best corrected visual acuity (BCVA) for each eye. Myopia and hyperopia were defined as minus and plus spherical equivalent refractive errors of greater than or equal to 0.50 DS respectively; astigmatism was defined as cylindrical correction of greater than or equal to 0.50 DC.

## Binocular vision examination for CI

Binocular vision investigation for CI included: measurement of near point of convergence (NPC) break and recovery using push-up method with RAF rule, measurement of distance and near lateral phoria using von Graefe phoria technique, smooth vergence testing to determine distance and near horizontal positive fusional vergence (PFV) amplitudes, and accommodative convergence over accommodation (AC/A) ratio using gradient method.<sup>12</sup> Binocular vision testing was done over BCVA refraction results.

## Criteria for diagnosing CI and ametropia

CI was diagnosed if three or more of the following clinical signs were present: a receded NPC (break point > 8 cm),<sup>3,12,19</sup> near decompensated exophoria greater than far exophoria (difference  $\geq 6^\Delta$ ),<sup>9,19</sup> low AC/A ratio (less than 3:1),<sup>3</sup> reduced PFV at near (<15 $^\Delta$  Sheard's criteria).<sup>9,19</sup> Ametropia was diagnosed if with BCVA refractive correction in place, participants presented normal results in binocular vision testing.<sup>20</sup>

## Statistical methods

Data was analyzed using IBM SPSS version 21 (SPSS, Inc., Chicago, IL, USA). Descriptive statistics such as frequencies, means with standard deviations, percentiles and 95% confidence intervals were computed for outcome variables. Normality of data was determined using the Shapiro Wilk p-values. Independent Sample Man Whitney U test was used to determine significant differences in age and CISS score among gender. Correlation between CISS score and clinical signs of CI were determined using Spearman's rho correlations. Welch ANOVA was used to test difference in CISS score between different ametropia. The sensitivity and specificity of the CISS to predict CI and ametropia were evaluated using

Receiver Operating Characteristics (ROC) curves. With analysis done within 95% confidence interval, a two-tailed p-value  $\leq 0.05$  was considered statistically significant.

## Results

The study enrolled 322 first year students out of which 22 were excluded with external ocular diseases ( $n = 9$ ), internal ocular diseases ( $n = 5$ ), strabismus ( $n = 3$ ), nystagmus ( $n = 1$ ), and on medical treatment ( $n = 4$ ). A total of 300 first year students comprising 167(55.7%) males and 133 (44.3%) females aged 18 to 27 years (mean = 21.58 (SD  $\pm 2.2$ ) years) were included in the study. Table 1 represents a descriptive data of all the parameters measured; some parameters were approximately normally distributed whilst others were not (Table 1). The minimum CISS score was 1 and the maximum was 42. The mean rank age for males (171.71) was significantly different from that (123.86) for the females (Mann–Whitney  $U$  test;  $p = 0.0001$ ). The mean rank CISS score for males (152.61) was not significantly different from that (147) for females (Mann–Whitney  $U$  test;  $p = 0.637$ ) Table 2 indicates the frequencies of refractive status, CI status and the number of clinical signs of convergence insufficiency observed among participants. The mean rank CISS score of participants with convergence insufficiency (254.02) was significantly higher than that (139.42) for those without convergence insufficiency (Mann–Whitney  $U$  test;  $p = 0.0001$ ). The descriptive measures for parameters of binocular vision for participants diagnosed with CI are indicated (Table 3). The NPC break and recovery of symptomatic CI participants were significantly higher than that for asymptomatic CI participants (Table 3). There were no significant differences in near exophoria, positive fusional vergence and AC/A ratio between symptomatic CI participants and asymptomatic CI participants (Table 3).

## Correlation between CISS score and signs of convergence insufficiency

There were significant correlations between CISS score and the clinical signs of CI (Table 4). Figs. 1, 2, 3 and 4 represent scatterplots of the association between specific parameters.

## Association between CISS score and number of clinical signs of CI

There was significant positive correlation between CISS score and the number of clinical signs of CI ( $r_s = 0.575$ ,  $p$ -value = 0.0001) as indicated (Fig. 5).

## The association between CI and Symptoms

Out of the 29 participants with convergence insufficiency, 26 were symptomatic and 3 were asymptomatic. Holding gender constant, symptomatic participants were more likely to experience convergence insufficiency (OR = 34.508, 95% CI = 9.945–119.688,  $p = 0.0001$ ) compared to asymptomatic

**Table 1** Descriptive measures of parameters investigated.

Parameter	25 <sup>th</sup> Percentile	Median	IQR	75 <sup>th</sup> Percentile	90 <sup>th</sup> Percentile	Mean $\pm$ SD	95% CI	Shapiro-Wilk p-values
CISS score	8	13	14	22	28.9	15.21 $\pm$ 10.02	14.07 – 16.35	0.0001
CISS males	7	15	15	22	27.2	15.26 $\pm$ 9.49	13.81 – 16.71	0.001
CISS females	8	12	14	22	31	15.16 $\pm$ 10.69	13.32 – 16.99	0.0001
Age males	20	22	4	24	25	22.13 $\pm$ 2.22	21.79 – 22.08	0.0001
Age females	19.5	21	3	22	24	20.90 $\pm$ 1.92	20.57 – 21.23	0.0001
NPC break	4	6	6	10	15	7.64 $\pm$ 4.476	7.13 – 8.15	0.0001
NPC rec	6	8	6	12	17	9.25 $\pm$ 4.787	8.71 – 9.80	0.0001
Exo N $\Delta$	2	3	1	4	4.8	3.09 $\pm$ 0.944	2.46 – 3.73	0.095
PFV blur N	18	20	3	21	24	19.91 $\pm$ 2.84	18 – 21.82	0.535
PFV break N	21	23	4	25	27.8	23.09 $\pm$ 2.95	21.11– 25.07	0.987
PFV rec N	16	18	3	19	21.6	17.73 $\pm$ 2.28	16.19 – 19.26	0.990
AC/A ratio	3	3	1	4	5.6	3.45 $\pm$ 1.04	2.79 – 4.15	0.011
Exo D $\Delta$	2	2	1	3	3	2.44 $\pm$ 0.796	2.21 – 2.67	0.0001
Eso D $\Delta$	1	2	1	2	2.8	1.73 $\pm$ 0.647	1.29 – 2.16	0.008
Eso N $\Delta$	2	3.5	4	6	6.9	3.80 $\pm$ 1.989	2.38 – 5.22	0.530
Myopia	0.50	0.75	0.5	1.00	1.75	0.89 $\pm$ 0.48	0.72 – 1.06	0.0001
Hyperopia	0.50	0.50	0.25	0.75	1.00	0.67 $\pm$ 0.31	0.59 – 0.74	0.0001
Astigmatism	0.50	0.63	0.25	0.75	1.25	0.75 $\pm$ .416	0.64 – 0.84	0.0001
CI (CISS)	23	33	16	38.5	43	30.24 $\pm$ 10.55	26.27 – 34.21	0.074
No CI (CISS)	7	12	12	19	25	13.61 $\pm$ 8.55	12.58 – 14.63	0.0001
CISS Ame	9	16	15	24	35	17.00 $\pm$ 9.49	15.56 – 18.44	0.0001
CISS Emm	5	11	13	18	30	12.13 $\pm$ 9.49	10.54 – 14.06	0.0001

IQR – interquartile range, CISS- convergence insufficiency symptom survey, Exo- exophoria, N- near, PFV- Positive fusional vergence, rec – recovery, Eso- esophoria, D- distance, CI- convergence insufficiency, Ame- Ametropia, Emm- Emmetropia.

**Table 2** Frequencies of various conditions investigated.

Condition	Frequency	Percentage (%)
<b>Refractive status</b>		
Emmetropia	114	38
Myopia	34	11.3
Hyperopia	62	20.7
Astigmatism	90	30
<b>Total</b>	300	100%
<b>Symptoms status</b>		
Symptomatic participants (CISS score $\geq$ 21)	84	28
Asymptomatic participants (CISS score < 21)	216	72
<b>Total</b>	300	100%
<b>Convergence Insufficiency status</b>		
Convergence insufficiency	29	9.7
No Convergence insufficiency	271	90.3
<b>Total</b>	300	100%
<b>Number of clinical signs of Convergence Insufficiency</b>		
0	183	61
1	53	17.7
2	35	11.7
3	16	5.33
4	13	4.33
<b>Total</b>	300	100%

participants (OR = 0.029, 95% CI = 0.01–0.101,  $p$  = 0.0001). There was a statistically significant ROC curve ( $p$  = 0.0001, AOC = 0.882, 95% CI = 0.804–0.960) with AOC indicating a good sensitivity and specificity of the CISS to predict CI (Fig. 6).

### The CISS score and ametropia

There was no significant correlation between CISS score, and magnitude of myopia ( $r_s$  = 0.002,  $p$ -value = 0.992), magnitude of hyperopia ( $r_s$  = 0.170,  $p$ -value = 0.188) and magnitude of astigmatism ( $r_s$  = -0.069,  $p$ -value = 0.521). The mean rank CISS score for participants with ametropia (166.64) was significantly different from that (124.16) for those with emmetropia (Mann–Whitney  $U$  test;  $p$  = 0.0001). Using Welch ANOVA, there was no significant difference in CISS score (Fig. 6) between the different types of ametropia ( $F$  (2,183) = 1.489,  $p$  = 0.228). There was a statistically significant ROC curve ( $p$  = 0.0001, AOC = 0.642, 95% CI = 0.577–0.706) with AOC indicating a poor sensitivity and specificity of the CISS to predict ametropia (Fig. 7).

### Discussion

The CISS score was found to be associated with the severity of specific CI signs and number of classic signs of CI among young adult university students. A higher CISS score among this young adult population is indicative of the severity of many signs of convergence insufficiency and vice versa. A

**Table 3** Descriptive parameters of participants with convergence insufficiency.

Parameter	25 <sup>th</sup> Percentile	Median	IQR	75 <sup>th</sup> Percentile	90 <sup>th</sup> Percentile	Mean $\pm$ SD	95% Confidence Interval	Shapiro-Wilk p-values
NPC break	12	15	6	18	20	15.03 $\pm$ 4.09	13.48 – 16.59	0.404
NPC rec	13.50	10	7	20.5	22	17.14 $\pm$ 4.25	15.52 - 18.75	0.566
Exo N $\Delta$	7	8	3	10	12	8.83 $\pm$ 2.07	8.04 - 9.62	0.003
PFV blur N	13	14	3	15.5	16	13.93 $\pm$ 2.65	12.92- 14.94	0.0001
PFV break N	16.5	18.0	4	20.0	22.0	17.93 $\pm$ 3.63	16.55-19.31	0.002
PFV rec N	11	14	5	16	17	13.28 $\pm$ 3.12	12.09-14.46	0.027
AC/A ratio	2	2	1	3	3	2.48 $\pm$ 0.34	2.24- 2.72	0.0001
Exo D $\Delta$	2	2	2	3.5	-	2.78 $\pm$ 1.39	1.71- 3.85	0.0001
Difference in parameters between symptomatic and asymptomatic convergence insufficiency participants								
	<b>Symptomatic CI</b>	<b>Asymptomatic CI</b>						
	Mean $\pm$ SD		t-value	P -values				
NPC break	15.62 $\pm$ 3.78	10.00 $\pm$ 2.00		2.442	0.021		0.90 – 10.33	
NPC recovery	17.81 $\pm$ 3.90	11.33 $\pm$ 2.52		2.784	0.010		1.70 – 11.25	
	Mean ranks		U	Z	P - values			
Exo N $\Delta$	15.6	9.83	23.5	-1.133	0.257			
PFV blur N	15.23	13.00	33.0	-0.439	0.663			
PFV break N	15.56	10.17	24.5	-1.061	0.289			
PFV rec N	15.31	12.33	31.0	-0.577	0.564			
AC/A ratio	14.64	18.00	30.0	-0.740	0.459			
Exo D $\Delta$	4.75	7.00	2	-0.920	0.357			

NPC – near point of convergence, IQR – interquartile range, Exo- exophoria, N- near, PFV- Positive fusional vergence, rec – recovery, D- distance, CI – convergence insufficiency

**Table 4** Correlation between CISS score and signs of CI.

Sign of CI	CISS score	
	Spearman Rho ( $r_s$ )	$p$ -value
NPC break	0.622	0.0001
NPC recovery	0.620	0.0001
Near exophoria	0.434	0.0001
PFV(blur at near)	- 0.359	0.0001
PFV (break at near)	- 0.306	0.0001
PFV (recovery at near)	- 0.326	0.0001
Gradient AC/A ratio	- 0.290	0.0001

higher CISS score (which directly predicts a greater likelihood of the presence of CI) was associated with all the signs of CI namely receded NPC break and recovery, higher decompensated exophoria at near, lower positive fusional vergence amplitudes and lower AC/A ratios. This result is not comparable with that found in children with three sign symptomatic convergence insufficiency.<sup>9</sup> The present study further emphasizes the difference in the relationship between symptoms and signs of CI in adults and children, as earlier indicated in difference in cut-off point for CISS.<sup>12,13</sup>


The linear relationships observed between the signs and symptoms of CI emphasize the point that treatment directed towards improving NPC, PFV, compensating for near exophoria and improving AC/A ratio goes a long way to treating this

condition as seen in the African case report.<sup>6</sup> The linear relationship between the CISS score, and the clinical signs of NPC break and near exophoria in this young adult population is inconsistent with that among symptomatic presbyopes with CI (for base line CISS scores)<sup>21</sup> and that among children 9 to 17 years old.<sup>9</sup>


The negative correlation between positive fusional vergence amplitudes and CISS score among the young adult population is not comparable to that for base line CISS scores in the study by Yi Pang et al.<sup>21</sup> Authors found no study that has compared AC/A ratio to CISS score. As high CISS score predict the presence of CI, low AC/A ratio is already known to be one of the clinical signs of CI.<sup>3</sup>

The present study results indicate that there is a greater likelihood for identifying symptomatic CI compared to asymptomatic CI using the CISS score. It also indicates that even though the chances are small, there is likelihood that asymptomatic participants (based on the CISS score) have CI. This indicates a limitation with the use of the CISS to screen for CI in young African adults; this observation was also emphasized by Horwood et al.<sup>11</sup> in their study among young adults. The diagnosis of CI should be based primarily on identifying the clinical signs rather than dependence on the CISS score. The CISS score when used in addition to clinical investigative testing for signs will give a definitive diagnosis of asymptomatic or symptomatic CI.


Of all the signs of CI investigated, only NPC was significantly higher among participants with symptomatic CI than

**Fig. 1** Positive correlation between CISS score and NPC break.





**Fig. 2** Positive correlation between CISS score and exophoria at near.



**Fig. 3** Negative correlation between CISS score and AC/A ratio.




Fig. 4 Scatterplot matrix showing negative correlation between CISS score and PFV amplitudes.

asymptomatic ones. The mean difference in NPC break (5.62 cm) is higher than the mean standard deviations of 3.78 cm and 2.00 cm respectively for the symptomatic and asymptomatic CI groups being compared. Also, the mean difference of 6.48 cm in NPC recovery is higher than the mean standard deviations of 3.90 cm and 2.52 cm respectively for the two groups. These differences in NPC break and recovery (between symptomatic and asymptomatic CI groups), therefore, are considered clinically meaningful. It can be inferred from this result that the sign of receded NPC may be accounting for the presence of significant symptoms among participants with CI. Symptomatic CI thus may result from the higher demand on convergence of the visual system during near focus. The CISS score cannot be used to predict ametropia within the population. This was further emphasized as participants with ametropia were asymptomatic and showed no differences per the CISS score (Table 2).

The validity and reliability of the CISS have been established in both children and adults.<sup>12,13</sup> A score of 21 or higher

differentiate adults with symptomatic CI from normal adults.<sup>12</sup> In our study, the average CISS score for all the subjects (15.21) suggest that averagely, participants were asymptomatic. The average CISS score (30.24) for participants diagnosed with CI is consistent with that of the CITT study group.<sup>9</sup> Generally, the diagnosis of CI in previous studies have been made using three<sup>9,22</sup> or four<sup>3,19</sup> of the clinical signs as employed in the present study. The prevalence of CI among the young adult Ghanaian university students (9.7%) is comparable to the 10% among young adult British university undergraduate and postgraduate students,<sup>11</sup> 7.7% among 65 Spanish university students (in a study where CI was diagnosed based on four clinical signs),<sup>23</sup> higher compared to 1.6% among Optometry students in Ghana<sup>2</sup> and lower compared to 29.6% among first year students in Nigeria.<sup>1</sup>

The study participants (university students) easily understood the concepts of blur and double vision and gave clear and reliable responses during testing. Authors, however, acknowledge as a limitation, the use of a less repeatable








Fig. 5 Positive correlation between CISS score and number of signs of CI.



Diagonal segments are produced by ties.

Fig. 6 ROC Curve indicating good sensitivity and specificity of CISS to predict CI. AOC is equal to 0.882.



Diagonal segments are produced by ties.

Fig. 7 ROC Curve indicating poor sensitivity and specificity of CISS to predict ametropia. AOU is equal to 0.642.

phoria measurement technique (von Graefe) in the present study, instead of a more repeatable technique like cover test.<sup>24</sup> von Graefe phoria technique produces greater heterophoria values than cover test among non-presbyopic patients.<sup>25</sup> It is thus possible that near phoria values in the present study may have been overestimated. The results in the present study are delimited to the procedures used.

## Conclusion

The CISS score is correlated with the severity and number of signs of CI and has a good sensitivity and specificity to predict CI in young adult Ghanaian students. The CISS poorly predicts the presence of ametropia among the young adult population. Its use in addition to clinical investigative testing gives a definitive diagnosis of symptomatic CI.

## Declaration of Competing Interest

The authors have no conflict of interest.

## References

- Ovenseri GO, Ovigwe PE. Vergence findings and horizontal vergence dysfunctions among first year university student in Benin City, Nigeria. *J Optom*. 2000. Available at: <https://doi.org/10.1016/j.optom.2016.01.004>.
- Darko-Takyi C, Owusu-Ansah A, Appiah-Eduenu C, Abu EK, Boadi-Kusi SB, Osei-Yaw A. Refractive and binocular vision status of optometry students, Ghana. *J Med Biomed Sci*. 2000. Available at: <https://doi.org/10.4314/jmbs.v5i2.4>.
- Scheiman M, Wick B. *Clinical management of binocular vision: heterophoric, accommodative and eye movement disorders*. 4th Ed. Philadelphia: J B Lippincott Williams & Wilkins; 2014.
- Barnhardt C, Cotter SA, Mitchell L, Scheiman M, Kulp MT. Symptoms in children with convergence insufficiency: before and after treatment. *Optom Vis Sci*. 2012;89:1512–1520.
- Convergence Insufficiency Treatment Trail Study Group. Randomized clinical trial of treatments for symptomatic convergence insufficiency in children. *Arch Ophthalmol*. 2008;126:1336–1349.
- Darko-Takyi C, Asiedu K. Symptomatic classic convergence insufficiency treated with dot card orthoptic exercise- a case report of an African patient. *Optom Reports*. 2016;5:5566.
- Rouse M, Borsting E, Mitchell G L, Cotter S A, Kulp M, Scheiman M, Barnhardt C, Bade A, Yamada T. Convergence Insufficiency Treatment Trial (CITT) Investigator Group. Validity of the convergence insufficiency symptom survey: a confirmatory study. *Optom Vis Sci*. 2009;86:357–363.
- Rouse MW, Hyman L, Hussein M, Solan H. Convergence insufficiency and reading study (CIRS) Group. Frequency of convergence insufficiency in optometry clinic settings. *Optom Vis Sci*. 1998;75:88–96.
- Bade A, Boas M, Gallaway M, Mitchell G L, Scheiman M, Kulp M T, Cotter S A, Rouse M. Relationship between clinical signs and symptoms of convergence insufficiency. *Optom Vis Sci*. 2013;90:988–995.
- Cohen Y, Segal O, Barkana Y, Lederman R, Zadok D, Pras E, Morad Y. Correlation between asthenopic symptoms and different measurements of convergence and reading comprehension and saccadic fixation eye movements. *Optometry*. 2010;81:28–34.
- Horwood A M, Toor S, Riddell P M. Screening for convergence insufficiency using CISS is not indicated in young adults. *Br J Ophthalmol*. 2000. Available at: [10.1136/bjophthalmol-2013-304533](https://doi.org/10.1136/bjophthalmol-2013-304533).
- Rouse MW, Borsting EJ, Mitchell GL, Scheiman M, Cotter SA, Cooper J, Kulp MT, London R, Wensveen J, Group CITT. Validity and reliability of the revised convergence insufficiency symptom survey in adults. *Ophthalmol. Physiol. Opt*. 2004;24:384–390.
- Borsting EJ, Rouse MW, Mitchell GL, Scheiman M, Cotter SA, Cooper J, Kulp MT, London R. Convergence insufficiency treatment trial group. Validity and reliability of the revised convergence insufficiency symptom survey in children aged 9-18 years. *Optom Vis Sci*. 2003;80:832–838.
- Horan LA, Ticho BH, Khammar AJ, Allen MS, Shah BA. Is The convergence insufficiency symptom survey specific for convergence insufficiency? a prospective, randomized study. *Am Orthopt J*. 2015;65:99–103.
- Borsting E, Rouse MW, Deland PN, Hovett S, Kimura D, Park M, Stephens B. Association of symptoms and convergence and accommodative insufficiency in school-age children. *Optometry*. 2003;74:25–34.
- Scheiman M, Gallaway M, Coulter R, Reinsteint F, Ciner E, Herzberg C, Parisi M. Prevalence of vision and ocular disease conditions in a clinical pediatric population. *J Am Optom Assoc*. 1996;67:193–202.
- Haine C. "The ophthalmic case historian,". In: Benjamin WJ, ed. *Borish's clinical refraction*. St. Louis MO, USA: Butterworth Heinemann; 2006. Ed.
- Benjamin WJ. *Borish's clinical refraction*. 2nd Ed. St. Louis, Missouri: Butterworth Heinemann Elsevier; 2006.
- Porcar E, Martinez-Palomera A. Prevalence of general dysfunctions in a population of university students. *Optom Vis Sci*. 1997;74:111–113.
- Lara F, Cacho P, García A, Megías R. General binocular disorders: prevalence in a clinic population. *Ophthalmic Physiol Opt*. 2001;21:70–74.
- Yi Pang Y, Teitelbaum B, Krall J. Factors associated with base-in prism treatment outcomes for convergence insufficiency in symptomatic presbyopes. *Clin Exp Optom*. 2012;95:192–197.
- Rouse MW, Borsting E, Hyman L, Mohammed H, Susan CA, Mary F, Scheiman M, Michael G, Paul D. The convergence insufficiency and reading study group. Frequency of convergence insufficiency among fifth and sixth graders. *Optom Vis Sci*. 1999;76:643–649.
- Porcar E, Martinez-Palomera A. Prevalence of general dysfunctions in a population of university students. *Optom Vis Sci*. 1997;74:111–113.
- Anstice NS, Davidson B, Field B, Mathanb J, Collins AV, Black JM. The repeatability and reproducibility of four techniques for measuring horizontal heterophoria: Implications for clinical practice. *J Optom*. 2020. <https://doi.org/10.1016/j.optom.2020.05.005>.
- Cantó-Cerdán M, Cacho-Martínez P, García-Muñoz Á. Measuring the heterophoria: agreement between two methods in non-presbyopic and presbyopic patients. *J Optom*. 2018;11:153–159.